

**CORRECTION TO
'DEFORMED CALABI–YAU COMPLETIONS'**

BERNHARD KELLER

The author is grateful to Dong Yang and Martin Kalck [1] for pointing out an error in Theorem 6.10 of [2]. To make the statement of the theorem correct, one has to assume in addition that the set of ‘minimal relations’ R generates the ideal I (from the definition of R , it only follows that R topologically generates the J -adic completion of I). Moreover, the condition 3) in the proof of the theorem should be replaced with

for all $n \geq 1$, the differential d maps V^{-n-1} to T_n and induces an isomorphism from V^{-n-1} onto the head of the $H^0(T_n)$ -bimodule $H^{-n}(T_n)$, where T_n denotes the dg category $T_{\mathcal{R}}(V^0 \oplus \cdots \oplus V^{-n})$.

REFERENCES

- [1] Martin Kalck and Dong Yang, electronic message, December 2012.
- [2] Bernhard Keller, *Deformed Calabi–Yau completions*, Journal für die reine und angewandte Mathematik (Crelles Journal) **654** (2011), 125–180, with an appendix by Michel Van den Bergh.

BERNHARD KELLER, U.F.R. DE MATHÉMATIQUES, INSTITUT DE MATHÉMATIQUES DE JUSSIEU,
U.M.R. 7586 DU CNRS, UNIVERSITÉ DENIS DIDEROT – PARIS 7, CASE 7012, 2 PLACE JUSSIEU,
75251 PARIS, CEDEX 05, FRANCE

E-mail address: `keller@math.jussieu.fr`