

Srinivasa Ramanujan

His life and his work

Michel Waldschmidt

Institut de Mathématiques de Jussieu & Paris VI

<http://www.imj-prg.fr/~michel.waldschmidt/>

Abstract

This lecture includes a few biographical informations about [Srinivasan Ramanujan](#). Among the topics which we discuss are [Euler](#) constant, nested roots, divergent series, [Ramanujan – Nagell](#) equation, partitions, [Ramanujan](#) tau function, [Hardy Littlewood](#) and the circle method, highly composite numbers and transcendence theory, the number π , and the lost notebook.

Srinivasa Ramanujan


Erode December 22, 1887 —

Chetput, (Madras), April 26, 1920



P.K. Srinivasan

(November 4, 1924-June 20, 2005)



PKS was the first biographer of Srinivas Ramanujan.

The Hindu, November 1, 2009
Passion for numbers by
Soudhamini

<http://beta.thehindu.com/education/article41732.ece>

Biography of Srinivasa Ramanujan

(December 22, 1887 — April 26, 1920)

1887 : born in Erode (near Tanjore)

1894-1903 : school in Kumbakonam

In 1900 he began to work on his own on mathematics summing geometric and arithmetic series.

Biography of Srinivasa Ramanujan

(December 22, 1887 — April 26, 1920)

1887 : born in Erode (near Tanjore)

1894-1903 : school in Kumbakonam

In 1900 he began to work on his own on mathematics summing geometric and arithmetic series.

Biography of Srinivasa Ramanujan

(December 22, 1887 — April 26, 1920)

1887 : born in Erode (near Tanjore)

1894-1903 : school in Kumbakonam

In 1900 he began to work on his own on mathematics summing geometric and arithmetic series.

Biography of Srinivasa Ramanujan

(December 22, 1887 — April 26, 1920)

1887 : born in Erode (near Tanjore)

1894-1903 : school in Kumbakonam

In 1900 he began to work on his own on mathematics summing geometric and arithmetic series.

Gopuram Sarangapani Kumbakonam



Sarangapani Sannidhi Street Kumbakonam



Ramanujan House Kumbakonam



Ramanujan House in Kumbakonam



Ramanujan House Kumbakonam



Town High School Kumbakonam



Town High School Kumbakonam

1903 : G.S.Carr - *A synopsis of elementary results — a book on pure mathematics* (1886) 5 000 formulae

$$\sqrt{x} + y = 7, \quad x + \sqrt{y} = 11$$

$$x = 9, \quad y = 4.$$

Town High School Kumbakonam

1903 : G.S.Carr - *A synopsis of elementary results — a book on pure mathematics* (1886) 5 000 formulae

$$\sqrt{x} + y = 7, \quad x + \sqrt{y} = 11$$

$$x = 9, \quad y = 4.$$

Town High School Kumbakonam

1903 : G.S.Carr - *A synopsis of elementary results — a book on pure mathematics* (1886) 5 000 formulae

$$\sqrt{x} + y = 7, \quad x + \sqrt{y} = 11$$

$$x = 9, \quad y = 4.$$

Biography (*continued*)

1903 (December) : exam at Madras University

1904 (January) : enters Government Arts College,
Kumbakonam

Sri K. Ranganatha Rao Prize

Subrahmanyam scholarship

Biography (*continued*)

1903 (December) : exam at Madras University

1904 (January) : enters Government Arts College,
Kumbakonam

Sri K. Ranganatha Rao Prize

Subrahmanyam scholarship

Biography (*continued*)

1903 (December) : exam at Madras University

1904 (January) : enters Government Arts College,
Kumbakonam

Sri K. Ranganatha Rao Prize

Subrahmanyam scholarship

MacTutor History of Mathematics

<http://www-history.mcs.st-andrews.ac.uk/>

By 1904 Ramanujan had begun to undertake deep research. He investigated the series

$$\sum_n \frac{1}{n}$$

and calculated Euler's constant to 15 decimal places.

He began to study the Bernoulli numbers, although this was entirely his own independent discovery.

MacTutor History of Mathematics

<http://www-history.mcs.st-andrews.ac.uk/>

By 1904 **Ramanujan** had begun to undertake deep research. He investigated the series

$$\sum_n \frac{1}{n}$$

and calculated **Euler's** constant to **15** decimal places.

He began to study the **Bernoulli** numbers, although this was entirely his own independent discovery.

Euler constant

$$S_N = \sum_{n=1}^N \frac{1}{n} = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots + \frac{1}{N}$$

$$\int_1^N \frac{dx}{x+1} < S_N < 1 + \int_1^N \frac{dx}{x}$$

$$\gamma = \lim_{N \rightarrow \infty} (S_N - \log N).$$

Euler constant

$$S_N = \sum_{n=1}^N \frac{1}{n} = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots + \frac{1}{N}$$

$$\int_1^N \frac{dx}{x+1} < S_N < 1 + \int_1^N \frac{dx}{x}$$

$$\gamma = \lim_{N \rightarrow \infty} (S_N - \log N).$$

Euler constant

$$S_N = \sum_{n=1}^N \frac{1}{n} = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots + \frac{1}{N}$$

$$\int_1^N \frac{dx}{x+1} < S_N < 1 + \int_1^N \frac{dx}{x}$$

$$\gamma = \lim_{N \rightarrow \infty} (S_N - \log N).$$

Reference



JEFFREY C. LAGARIAS

*Euler's constant : Euler's work
and modern developments*

Bulletin Amer. Math. Soc. **50**
(2013), No. 4, 527–628.

[arXiv:1303.1856](https://arxiv.org/abs/1303.1856) [math.NT]

Bibliography : 314 references.

Euler archives and Eneström index

The Euler Archive

A digital library dedicated to the work and life of Leonhard Euler



<http://eulerarchive.maa.org/>

Gustaf Eneström (1852–1923)

*Die Schriften Euler's
chronologisch nach den Jahren
geordnet, in denen sie verfasst
worden sind*

Jahresbericht der Deutschen
Mathematiker-Vereinigung,
1913.



Gustaf Eneström.
Efter fotografi.

<http://www.math.dartmouth.edu/~euler/index/enestrom.html>

<http://www.eulerarchive.org/>



The screenshot shows the Euler Archive website. At the top, the word "eulerarchive" is written in a large, red, italicized font. Below this is a navigation bar with four tabs: "Home", "About", "News", and "Race". The main content area features a large image of a yellow Formula 1 race car. Overlaid on the left side of the car is a text box with Thai text. The text reads: "F1: รถใหม่ ของเฟอร์รารี ที่เกือบจะ สมบูรณ์" (F1: Ferrari's new car that is almost perfect). Below this is a paragraph of text: "แม้ว่าทีม 2014 นั้น สามารถ ก่อตั้งเป็นทางการ ในการเปิดตัว เมื่อ วันที่ 15 มีนาคมที่เปิดตัวอย่างเป็นทางการ ที่ เฟอร์รารี แต่ มี พยายามที่จะ ยืดเยื้อ ส่งเสริมสมรรถนะ ในการ โต้ตอบของรถ ที่สมบูรณ์ ตามที่ มาของวิศวกร เฟอร์รารี โดยอ้างว่าส่วนหัว ใหม่นี้ ครอบ ครอบ (fullrod) เป็นรุ่น 2012-2013 กลับเป็น 2011 แล้วบน ก้าน ครอบที่ นั้น". At the bottom left of the text box is a red button with the text "VIEW MORE".

(Référence [86] of the text by Lagarias)

Harmonic numbers

$$H_1 = 1, \quad H_2 = 1 + \frac{1}{2} = \frac{3}{2}, \quad H_3 = 1 + \frac{1}{2} + \frac{1}{3} = \frac{11}{6},$$

$$H_n = 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} = \sum_{j=1}^n \frac{1}{j}.$$

Sequence :

$$1, \quad \frac{3}{2}, \quad \frac{11}{6}, \quad \frac{25}{12}, \quad \frac{137}{60}, \quad \frac{49}{20}, \quad \frac{363}{140}, \quad \frac{761}{280}, \quad \frac{7129}{2520}, \dots$$

Harmonic numbers

$$H_1 = 1, \quad H_2 = 1 + \frac{1}{2} = \frac{3}{2}, \quad H_3 = 1 + \frac{1}{2} + \frac{1}{3} = \frac{11}{6},$$

$$H_n = 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} = \sum_{j=1}^n \frac{1}{j}.$$

Sequence :

$$1, \quad \frac{3}{2}, \quad \frac{11}{6}, \quad \frac{25}{12}, \quad \frac{137}{60}, \quad \frac{49}{20}, \quad \frac{363}{140}, \quad \frac{761}{280}, \quad \frac{7129}{2520}, \dots$$

Harmonic numbers

$$H_1 = 1, \quad H_2 = 1 + \frac{1}{2} = \frac{3}{2}, \quad H_3 = 1 + \frac{1}{2} + \frac{1}{3} = \frac{11}{6},$$

$$H_n = 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} = \sum_{j=1}^n \frac{1}{j}.$$

Sequence :

$$1, \quad \frac{3}{2}, \quad \frac{11}{6}, \quad \frac{25}{12}, \quad \frac{137}{60}, \quad \frac{49}{20}, \quad \frac{363}{140}, \quad \frac{761}{280}, \quad \frac{7129}{2520}, \dots$$

The online encyclopaedia of integer sequences

<https://oeis.org/>

Neil J. A. Sloane



Numerators and denominators

Numerators : <https://oeis.org/A001008>

1, 3, 11, 25, 137, 49, 363, 761, 7129, 7381, 83711, 86021, 1145993,
1171733, 1195757, 2436559, 42142223, 14274301, 275295799,
55835135, 18858053, 19093197, 444316699, 1347822955, ...

Denominators : <https://oeis.org/A002805>

1, 2, 6, 12, 60, 20, 140, 280, 2520, 2520, 27720, 27720, 360360,
360360, 360360, 720720, 12252240, 4084080, 77597520,
15519504, 5173168, 5173168, 118982864, 356948592, ...

Euler (1731)

De progressionibus harmonicis observationes

The sequence

$$H_n - \log n$$

has a limit $\gamma = 0,577\,218\dots$
when n tends to infinity.

Leonhard Euler
(1707–1783)



Moreover,

$$\gamma = \sum_{m=2}^{\infty} (-1)^m \frac{\zeta(m)}{m}.$$

Riemann zeta function



$$\begin{aligned}\zeta(s) &= \sum_{n \geq 1} \frac{1}{n^s} \\ &= \prod_p \frac{1}{1 - p^{-s}}\end{aligned}$$



Euler : $s \in \mathbb{R}$.

Riemann : $s \in \mathbb{C}$.

Numerical value of Euler's constant

The online encyclopaedia of integer sequences

<https://oeis.org/A001620>

Decimal expansion of Euler's constant
(or Euler–Mascheroni constant) gamma.

Yee (2010) computed 29 844 489 545 decimal digits of gamma.

$\gamma = 0,577\ 215\ 664\ 901\ 532\ 860\ 606\ 512\ 090\ 082\ 402\ 431\ 042\ \dots$

Euler constant

Euler–Mascheroni constant

$$\gamma = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} - \log n \right) = 0.577\,215\,664\,9\dots$$



Neil J. A. Sloane's encyclopaedia

<http://www.research.att.com/~njas/sequences/A001620>

Bernoulli numbers



$$B_0 = 1, \quad \sum_{k=0}^{n-1} \binom{n}{k} B_k = 0 \quad \text{for } n > 1.$$

Jacob Bernoulli (1654 – 1705)

$$B_0 + 2B_1 = 0 \qquad B_1 = -\frac{1}{2}$$

$$B_0 + 3B_1 + 3B_2 = 0 \qquad B_2 = \frac{1}{6}$$

$$B_0 + 4B_1 + 6B_2 + 4B_3 = 0 \qquad B_3 = 0$$

$$B_0 + 5B_1 + 10B_2 + 10B_3 + 5B_4 = 0 \qquad B_4 = -\frac{1}{30}$$

⋮

Bernoulli numbers



$$B_0 = 1, \quad \sum_{k=0}^{n-1} \binom{n}{k} B_k = 0 \quad \text{for } n > 1.$$

Jacob Bernoulli (1654 – 1705)

$$B_0 + 2B_1 = 0 \qquad B_1 = -\frac{1}{2}$$

$$B_0 + 3B_1 + 3B_2 = 0 \qquad B_2 = \frac{1}{6}$$

$$B_0 + 4B_1 + 6B_2 + 4B_3 = 0 \qquad B_3 = 0$$

$$B_0 + 5B_1 + 10B_2 + 10B_3 + 5B_4 = 0 \qquad B_4 = -\frac{1}{30}$$

⋮

Kumbakonam

1905 : Fails final exam

1906 : Enters Pachaiyappa's College, Madras
III, goes back to Kumbakonam

1907 (December) : Fails final exam.

1908 : continued fractions and divergent series

1909 (April) : underwent an operation

1909 (July 14) : marriage with S Janaki Ammal (1900—1994)

Kumbakonam

1905 : Fails final exam

1906 : Enters Pachaiyappa's College, Madras

III, goes back to Kumbakonam

1907 (December) : Fails final exam.

1908 : continued fractions and divergent series

1909 (April) : underwent an operation

1909 (July 14) : marriage with S Janaki Ammal (1900—1994)

Kumbakonam

1905 : Fails final exam

1906 : Enters Pachaiyappa's College, Madras

III, goes back to Kumbakonam

1907 (December) : Fails final exam.

1908 : continued fractions and divergent series

1909 (April) : underwent an operation

1909 (July 14) : marriage with S Janaki Ammal (1900—1994)

Kumbakonam

1905 : Fails final exam

1906 : Enters Pachaiyappa's College, Madras
III, goes back to Kumbakonam

1907 (December) : Fails final exam.

1908 : **continued fractions** and divergent series

1909 (April) : underwent an operation

1909 (July 14) : marriage with **S Janaki Ammal** (1900—1994)

Kumbakonam

1905 : Fails final exam

1906 : Enters Pachaiyappa's College, Madras
III, goes back to Kumbakonam

1907 (December) : Fails final exam.

1908 : **continued fractions** and divergent series

1909 (April) : underwent an operation

1909 (July 14) : marriage with **S Janaki Ammal** (1900—1994)

Kumbakonam

1905 : Fails final exam

1906 : Enters Pachaiyappa's College, Madras
III, goes back to Kumbakonam

1907 (December) : Fails final exam.

1908 : **continued fractions** and divergent series

1909 (April) : underwent an operation

1909 (July 14) : marriage with **S Janaki Ammal** (1900—1994)

S Janaki Ammal



Madras

1910 : meets [Ramaswami Aiyar](#)

1911 : first mathematical paper

1912 : clerk office, Madras Port Trust — [Sir Francis Spring](#) and [Sir Gilbert Walker](#) get a scholarship for him from the University of Madras starting May 1913 for 2 years.

Madras

1910 : meets [Ramaswami Aiyar](#)

1911 : first mathematical paper

1912 : clerk office, Madras Port Trust — [Sir Francis Spring](#) and [Sir Gilbert Walker](#) get a scholarship for him from the University of Madras starting May 1913 for 2 years.

Madras

1910 : meets [Ramaswami Aiyar](#)

1911 : first mathematical paper

1912 : clerk office, Madras Port Trust — [Sir Francis Spring](#) and [Sir Gilbert Walker](#) get a scholarship for him from the University of Madras starting May 1913 for 2 years.

1912 Questions in the Journal of the Indian Mathematical Society

$$\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \cdots}}}} = ?$$

$$\sqrt{6 + 2\sqrt{7 + 3\sqrt{8 + 4\sqrt{9 + \cdots}}}} = ?$$

1912 Questions in the Journal of the Indian Mathematical Society

$$\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \cdots}}}} = ?$$

$$\sqrt{6 + 2\sqrt{7 + 3\sqrt{8 + 4\sqrt{9 + \cdots}}}} = ?$$

Answers from Ramanujan

$$\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \cdots}}}} = 3$$

$$\sqrt{6 + 2\sqrt{7 + 3\sqrt{8 + 4\sqrt{9 + \cdots}}}} = 4$$

Answers from Ramanujan

$$\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \cdots}}}} = 3$$

$$\sqrt{6 + 2\sqrt{7 + 3\sqrt{8 + 4\sqrt{9 + \cdots}}}} = 4$$

“Proofs” $n(n + 2)$

$$(n + 2)^2 = 1 + (n + 1)(n + 3)$$

$$n(n + 2) = n\sqrt{1 + (n + 1)(n + 3)}$$

$$f(n) = n(n + 2)$$

$$f(n) = n\sqrt{1 + f(n + 1)}$$

$$f(n) = n\sqrt{1 + (n + 1)\sqrt{1 + f(n + 2)}}$$

$$= n\sqrt{1 + (n + 1)\sqrt{1 + (n + 2)\sqrt{1 + (n + 3)\cdots}}}$$

$$f(1) = 3$$

“Proofs” $n(n + 2)$

$$(n + 2)^2 = 1 + (n + 1)(n + 3)$$

$$n(n + 2) = n\sqrt{1 + (n + 1)(n + 3)}$$

$$f(n) = n(n + 2)$$

$$f(n) = n\sqrt{1 + f(n + 1)}$$

$$f(n) = n\sqrt{1 + (n + 1)\sqrt{1 + f(n + 2)}}$$

$$= n\sqrt{1 + (n + 1)\sqrt{1 + (n + 2)\sqrt{1 + (n + 3)\cdots}}}$$

$$f(1) = 3$$

“Proofs” $n(n + 2)$

$$(n + 2)^2 = 1 + (n + 1)(n + 3)$$

$$n(n + 2) = n\sqrt{1 + (n + 1)(n + 3)}$$

$$f(n) = n(n + 2)$$

$$f(n) = n\sqrt{1 + f(n + 1)}$$

$$f(n) = n\sqrt{1 + (n + 1)\sqrt{1 + f(n + 2)}}$$

$$= n\sqrt{1 + (n + 1)\sqrt{1 + (n + 2)\sqrt{1 + (n + 3)\cdots}}}$$

$$f(1) = 3$$

“Proofs” $n(n + 2)$

$$(n + 2)^2 = 1 + (n + 1)(n + 3)$$

$$n(n + 2) = n\sqrt{1 + (n + 1)(n + 3)}$$

$$f(n) = n(n + 2)$$

$$f(n) = n\sqrt{1 + f(n + 1)}$$

$$f(n) = n\sqrt{1 + (n + 1)\sqrt{1 + f(n + 2)}}$$

$$= n\sqrt{1 + (n + 1)\sqrt{1 + (n + 2)\sqrt{1 + (n + 3)\cdots}}}$$

$$f(1) = 3$$

“Proofs” $n(n + 2)$

$$(n + 2)^2 = 1 + (n + 1)(n + 3)$$

$$n(n + 2) = n\sqrt{1 + (n + 1)(n + 3)}$$

$$f(n) = n(n + 2)$$

$$f(n) = n\sqrt{1 + f(n + 1)}$$

$$f(n) = n\sqrt{1 + (n + 1)\sqrt{1 + f(n + 2)}}$$

$$= n\sqrt{1 + (n + 1)\sqrt{1 + (n + 2)\sqrt{1 + (n + 3)\cdots}}}$$

$$f(1) = 3$$

“Proofs” $n(n + 2)$

$$(n + 2)^2 = 1 + (n + 1)(n + 3)$$

$$n(n + 2) = n\sqrt{1 + (n + 1)(n + 3)}$$

$$f(n) = n(n + 2)$$

$$f(n) = n\sqrt{1 + f(n + 1)}$$

$$f(n) = n\sqrt{1 + (n + 1)\sqrt{1 + f(n + 2)}}$$

$$= n\sqrt{1 + (n + 1)\sqrt{1 + (n + 2)\sqrt{1 + (n + 3)\cdots}}}$$

$$f(1) = 3$$

“Proofs” $n(n + 2)$

$$(n + 2)^2 = 1 + (n + 1)(n + 3)$$

$$n(n + 2) = n\sqrt{1 + (n + 1)(n + 3)}$$

$$f(n) = n(n + 2)$$

$$f(n) = n\sqrt{1 + f(n + 1)}$$

$$f(n) = n\sqrt{1 + (n + 1)\sqrt{1 + f(n + 2)}}$$

$$= n\sqrt{1 + (n + 1)\sqrt{1 + (n + 2)\sqrt{1 + (n + 3)\cdots}}}$$

$$f(1) = 3$$

“Proofs” $n(n + 3)$

$$(n + 3)^2 = n + 5 + (n + 1)(n + 4)$$

$$n(n + 3) = n\sqrt{n + 5 + (n + 1)(n + 4)}$$

$$g(n) = n(n + 3)$$

$$g(n) = n\sqrt{n + 5 + g(n + 1)}$$

$$g(n) = n\sqrt{n + 5 + (n + 1)\sqrt{n + 6 + g(n + 2)}}$$

$$= n\sqrt{n + 5 + (n + 1)\sqrt{n + 6 + (n + 2)\sqrt{n + 7 + \dots}}}$$

$$g(1) = 4$$

“Proofs” $n(n + 3)$

$$(n + 3)^2 = n + 5 + (n + 1)(n + 4)$$

$$n(n + 3) = n\sqrt{n + 5 + (n + 1)(n + 4)}$$

$$g(n) = n(n + 3)$$

$$g(n) = n\sqrt{n + 5 + g(n + 1)}$$

$$g(n) = n\sqrt{n + 5 + (n + 1)\sqrt{n + 6 + g(n + 2)}}$$

$$= n\sqrt{n + 5 + (n + 1)\sqrt{n + 6 + (n + 2)\sqrt{n + 7 + \dots}}}$$

$$g(1) = 4$$

“Proofs” $n(n + 3)$

$$(n + 3)^2 = n + 5 + (n + 1)(n + 4)$$

$$n(n + 3) = n\sqrt{n + 5 + (n + 1)(n + 4)}$$

$$g(n) = n(n + 3)$$

$$g(n) = n\sqrt{n + 5 + g(n + 1)}$$

$$g(n) = n\sqrt{n + 5 + (n + 1)\sqrt{n + 6 + g(n + 2)}}$$

$$= n\sqrt{n + 5 + (n + 1)\sqrt{n + 6 + (n + 2)\sqrt{n + 7 + \dots}}}$$

$$g(1) = 4$$

“Proofs” $n(n + 3)$

$$(n + 3)^2 = n + 5 + (n + 1)(n + 4)$$

$$n(n + 3) = n\sqrt{n + 5 + (n + 1)(n + 4)}$$

$$g(n) = n(n + 3)$$

$$g(n) = n\sqrt{n + 5 + g(n + 1)}$$

$$g(n) = n\sqrt{n + 5 + (n + 1)\sqrt{n + 6 + g(n + 2)}}$$

$$= n\sqrt{n + 5 + (n + 1)\sqrt{n + 6 + (n + 2)\sqrt{n + 7 + \dots}}}$$

$$g(1) = 4$$

“Proofs” $n(n + 3)$

$$(n + 3)^2 = n + 5 + (n + 1)(n + 4)$$

$$n(n + 3) = n\sqrt{n + 5 + (n + 1)(n + 4)}$$

$$g(n) = n(n + 3)$$

$$g(n) = n\sqrt{n + 5 + g(n + 1)}$$

$$g(n) = n\sqrt{n + 5 + (n + 1)\sqrt{n + 6 + g(n + 2)}}$$

$$= n\sqrt{n + 5 + (n + 1)\sqrt{n + 6 + (n + 2)\sqrt{n + 7 + \dots}}}$$

$$g(1) = 4$$

“Proofs” $n(n + 3)$

$$(n + 3)^2 = n + 5 + (n + 1)(n + 4)$$

$$n(n + 3) = n\sqrt{n + 5 + (n + 1)(n + 4)}$$

$$g(n) = n(n + 3)$$

$$g(n) = n\sqrt{n + 5 + g(n + 1)}$$

$$g(n) = n\sqrt{n + 5 + (n + 1)\sqrt{n + 6 + g(n + 2)}}$$

$$= n\sqrt{n + 5 + (n + 1)\sqrt{n + 6 + (n + 2)\sqrt{n + 7 + \dots}}}$$

$$g(1) = 4$$

“Proofs” $n(n + 3)$

$$(n + 3)^2 = n + 5 + (n + 1)(n + 4)$$

$$n(n + 3) = n\sqrt{n + 5 + (n + 1)(n + 4)}$$

$$g(n) = n(n + 3)$$

$$g(n) = n\sqrt{n + 5 + g(n + 1)}$$

$$g(n) = n\sqrt{n + 5 + (n + 1)\sqrt{n + 6 + g(n + 2)}}$$

$$= n\sqrt{n + 5 + (n + 1)\sqrt{n + 6 + (n + 2)\sqrt{n + 7 + \dots}}}$$

$$g(1) = 4$$

Letter of S. Ramanujan to M.J.M. Hill in 1912

$$1 + 2 + 3 + \cdots + \infty = -\frac{1}{12}$$

$$1^2 + 2^2 + 3^2 + \cdots + \infty^2 = 0$$

$$1^3 + 2^3 + 3^3 + \cdots + \infty^3 = \frac{1}{120}$$



Letter of S. Ramanujan to M.J.M. Hill in 1912

$$1 + 2 + 3 + \cdots + \infty = -\frac{1}{12}$$

$$1^2 + 2^2 + 3^2 + \cdots + \infty^2 = 0$$

$$1^3 + 2^3 + 3^3 + \cdots + \infty^3 = \frac{1}{120}$$



Letter of S. Ramanujan to M.J.M. Hill in 1912

$$1 + 2 + 3 + \cdots + \infty = -\frac{1}{12}$$

$$1^2 + 2^2 + 3^2 + \cdots + \infty^2 = 0$$

$$1^3 + 2^3 + 3^3 + \cdots + \infty^3 = \frac{1}{120}$$



Letter of S. Ramanujan to M.J.M. Hill in 1912

$$1 + 2 + 3 + \cdots + \infty = -\frac{1}{12}$$

$$1^2 + 2^2 + 3^2 + \cdots + \infty^2 = 0$$

$$1^3 + 2^3 + 3^3 + \cdots + \infty^3 = \frac{1}{120}$$



Answer of M.J.M. Hill in 1912

$$1 + 2 + 3 + \cdots + n = \frac{1}{2}n(n + 1)$$

$$1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{n(2n + 1)(n + 1)}{6}$$

$$1^3 + 2^3 + 3^3 + \cdots + n^3 = \left(\frac{n(n + 1)}{2} \right)^2$$

Renormalisation of divergent series



Leonhard Euler

(1707 – 1783)

Introductio in analysin infinitorum

(1748)

Euler

Values of **Riemann** zeta function at negative integers :

$$\zeta(-k) = -\frac{B_{k+1}}{k+1} \quad (n \geq 1)$$

$$\zeta(-2n) = 1^{2n} + 2^{2n} + 3^{2n} + 4^{2n} + \dots = 0 \quad (n \geq 1)$$

$$\zeta(-1) = 1 + 2 + 3 + 4 + \dots = -\frac{1}{12}$$

$$\zeta(-3) = 1^3 + 2^3 + 3^3 + 4^3 + \dots = \frac{1}{120}$$

$$\zeta(-5) = 1^5 + 2^5 + 3^5 + 4^5 + \dots = -\frac{1}{252}$$

Euler

Values of **Riemann** zeta function at negative integers :

$$\zeta(-k) = -\frac{B_{k+1}}{k+1} \quad (n \geq 1)$$

$$\zeta(-2n) = 1^{2n} + 2^{2n} + 3^{2n} + 4^{2n} + \dots = 0 \quad (n \geq 1)$$

$$\zeta(-1) = 1 + 2 + 3 + 4 + \dots = -\frac{1}{12}$$

$$\zeta(-3) = 1^3 + 2^3 + 3^3 + 4^3 + \dots = \frac{1}{120}$$

$$\zeta(-5) = 1^5 + 2^5 + 3^5 + 4^5 + \dots = -\frac{1}{252}$$

Euler

Values of **Riemann** zeta function at negative integers :

$$\zeta(-k) = -\frac{B_{k+1}}{k+1} \quad (n \geq 1)$$

$$\zeta(-2n) = 1^{2n} + 2^{2n} + 3^{2n} + 4^{2n} + \dots = 0 \quad (n \geq 1)$$

$$\zeta(-1) = 1 + 2 + 3 + 4 + \dots = -\frac{1}{12}$$

$$\zeta(-3) = 1^3 + 2^3 + 3^3 + 4^3 + \dots = \frac{1}{120}$$

$$\zeta(-5) = 1^5 + 2^5 + 3^5 + 4^5 + \dots = -\frac{1}{252}$$

Euler

Values of **Riemann** zeta function at negative integers :

$$\zeta(-k) = -\frac{B_{k+1}}{k+1} \quad (n \geq 1)$$

$$\zeta(-2n) = 1^{2n} + 2^{2n} + 3^{2n} + 4^{2n} + \dots = 0 \quad (n \geq 1)$$

$$\zeta(-1) = 1 + 2 + 3 + 4 + \dots = -\frac{1}{12}$$

$$\zeta(-3) = 1^3 + 2^3 + 3^3 + 4^3 + \dots = \frac{1}{120}$$

$$\zeta(-5) = 1^5 + 2^5 + 3^5 + 4^5 + \dots = -\frac{1}{252}$$

G.H. Hardy : Divergent Series (1949)



Niels Henrik Abel
(1802 – 1829)

Divergent series are the invention of the devil, and it is shameful to base on them any demonstration whatsoever.

Letters to H.F. Baker and E.W. Hobson in 1912

No answer to his letters to H.F. Baker and E.W. Hobson in 1912...

Letter of Ramanujan to Hardy

(January 16, 1913)

I have had no university education but I have undergone the ordinary school course. After leaving school I have been employing the spare time at my disposal to work at mathematics. I have not trodden through the conventional regular course which is followed in a university course, but I am striking out a new path for myself. I have made a special investigation of divergent series in general and the results I get are termed by the local mathematicians as “startling”.

Godfrey Harold Hardy (1877 – 1947)



John Edensor Littlewood (1885 – 1977)



Hardy and Littlewood



Letter from Ramanujan to Hardy

(January 16, 1913)

$$1 - 2 + 3 - 4 + \cdots = \frac{1}{4}$$

$$1 - 1! + 2! - 3! + \cdots = .596 \cdots$$

Answer from Hardy

(February 8, 1913)

I was exceedingly interested by your letter and by the theorems which you state. You will however understand that, before I can judge properly of the value of what you have done, it is essential that I should see proofs of some of your assertions. Your results seem to me to fall into roughly three classes :

(1) there are a number of results that are already known, or easily deducible from known theorems ;

(2) there are results which, so far as I know, are new and interesting, but interesting rather from their curiosity and apparent difficulty than their importance ;

(3) there are results which appear to be new and important. . .

Answer from Hardy

(February 8, 1913)

I was exceedingly interested by your letter and by the theorems which you state. You will however understand that, before I can judge properly of the value of what you have done, it is essential that I should see proofs of some of your assertions. Your results seem to me to fall into roughly three classes :

(1) there are a number of results that are already known, or easily deducible from known theorems ;

(2) there are results which, so far as I know, are new and interesting, but interesting rather from their curiosity and apparent difficulty than their importance ;

(3) there are results which appear to be new and important. . .

Answer from Hardy

(February 8, 1913)

I was exceedingly interested by your letter and by the theorems which you state. You will however understand that, before I can judge properly of the value of what you have done, it is essential that I should see proofs of some of your assertions. Your results seem to me to fall into roughly three classes :

(1) there are a number of results that are already known, or easily deducible from known theorems ;

(2) there are results which, so far as I know, are new and interesting, but interesting rather from their curiosity and apparent difficulty than their importance ;

(3) there are results which appear to be new and important. . .

Answer from Hardy

(February 8, 1913)

I was exceedingly interested by your letter and by the theorems which you state. You will however understand that, before I can judge properly of the value of what you have done, it is essential that I should see proofs of some of your assertions. Your results seem to me to fall into roughly three classes :

(1) there are a number of results that are already known, or easily deducible from known theorems;

(2) there are results which, so far as I know, are new and interesting, but interesting rather from their curiosity and apparent difficulty than their importance;

(3) there are results which appear to be new and important. . .

1913–1920

1913, February 27 : New letter from Ramanujan to Hardy

1913 : Visit of Neville to India

1914, March 17 to April 14 : travel to Cambridge.

1918 : (May) Fellow of the Royal Society
(November) Fellow of Trinity College, Cambridge.

1919, February 27 to March 13 : travel back to India.

1913–1920

1913, February 27 : New letter from Ramanujan to Hardy

1913 : Visit of Neville to India

1914, March 17 to April 14 : travel to Cambridge.

1918 : (May) Fellow of the Royal Society
(November) Fellow of Trinity College, Cambridge.

1919, February 27 to March 13 : travel back to India.

1913–1920

1913, February 27 : New letter from Ramanujan to Hardy

1913 : Visit of Neville to India

1914, March 17 to April 14 : travel to Cambridge.

1918 : (May) Fellow of the Royal Society
(November) Fellow of Trinity College, Cambridge.

1919, February 27 to March 13 : travel back to India.

1913–1920

1913, February 27 : New letter from Ramanujan to Hardy

1913 : Visit of Neville to India

1914, March 17 to April 14 : travel to Cambridge.

1918 : (May) Fellow of the Royal Society
(November) Fellow of Trinity College, Cambridge.

1919, February 27 to March 13 : travel back to India.

1913–1920

1913, February 27 : New letter from Ramanujan to Hardy

1913 : Visit of Neville to India

1914, March 17 to April 14 : travel to Cambridge.

1918 : (May) Fellow of the Royal Society
(November) Fellow of Trinity College, Cambridge.

1919, February 27 to March 13 : travel back to India.

Ramanujan – Taxi Cab Number 1729

Hardy's obituary of Ramanujan :

I had ridden in taxi-cab No 1729, and remarked that the number $(7 \cdot 13 \cdot 19)$ seemed to me a rather dull one. . .

$$1729 = 1^3 + 12^3 = 9^3 + 10^3$$

$$12^3 = 1728, \quad 9^3 = 729$$

Ramanujan – Taxi Cab Number 1729

Hardy's obituary of Ramanujan :

I had ridden in taxi-cab No 1729, and remarked that the number $(7 \cdot 13 \cdot 19)$ seemed to me a rather dull one. . .

$$1729 = 1^3 + 12^3 = 9^3 + 10^3$$

$$12^3 = 1728, \quad 9^3 = 729$$

Ramanujan – Taxi Cab Number 1729

Hardy's obituary of Ramanujan :

I had ridden in taxi-cab No 1729, and remarked that the number $(7 \cdot 13 \cdot 19)$ seemed to me a rather dull one. . .

$$1729 = 1^3 + 12^3 = 9^3 + 10^3$$

$$12^3 = 1728, \quad 9^3 = 729$$

Narendra Jadhav — Taxi Cab Number 1729

Narendra Jadhav (born 1953) is a noted Indian bureaucrat, economist, social scientist, writer and educationist. He is a member of Planning Commission of India as well as a member of National Advisory Council (NAC), since 31 May 2010. Prior to this, he had worked with International Monetary Fund (IMF) and headed economic research at Reserve Bank of India (RBI).



He was Vice-Chancellor (from 24 August 2006 to 15 June 2009) of University of Pune
Author of *Outcaste – A Memoir, Life and Triumphs of an Untouchable Family In India* (2003).

<http://www.drnarendrajadhav.info>

Ramanujan – Taxi Cab Number 1729

$$12^3 = 1728, \quad 9^3 = 729$$

$$50 = 7^2 + 1^2 = 5^2 + 5^2$$

$$\begin{aligned} 4\,104 &= 2^3 + 16^3 = 9^3 + 15^3 \\ 13\,832 &= 2^3 + 24^3 = 18^3 + 20^3 \\ 40\,033 &= 9^3 + 34^3 = 16^3 + 33^3 \\ &\vdots \end{aligned}$$

Ramanujan – Taxi Cab Number 1729

$$12^3 = 1728, \quad 9^3 = 729$$

$$50 = 7^2 + 1^2 = 5^2 + 5^2$$

$$\begin{aligned} 4\,104 &= 2^3 + 16^3 = 9^3 + 15^3 \\ 13\,832 &= 2^3 + 24^3 = 18^3 + 20^3 \\ 40\,033 &= 9^3 + 34^3 = 16^3 + 33^3 \\ &\vdots \end{aligned}$$

Leonhard Euler (1707 – 1783)



$$59^4 + 158^4 = 133^4 + 134^4 = 635\,318\,657$$

Diophantine equations

$$x^3 + y^3 + z^3 = w^3$$

$$(x, y, z, w) = (3, 4, 5, 6)$$

$$3^3 + 4^3 + 5^3 = 27 + 64 + 125 = 216 = 6^3$$

Parametric solution :

$$x = 3a^2 + 5ab - 5b^2$$

$$z = 5a^2 - 5ab - 3b^2$$

$$y = 4a^2 - 4ab + 6b^2$$

$$w = 6a^2 - 4ab + 4b^2$$

Diophantine equations

$$x^3 + y^3 + z^3 = w^3$$

$$(x, y, z, w) = (3, 4, 5, 6)$$

$$3^3 + 4^3 + 5^3 = 27 + 64 + 125 = 216 = 6^3$$

Parametric solution :

$$x = 3a^2 + 5ab - 5b^2$$

$$z = 5a^2 - 5ab - 3b^2$$

$$y = 4a^2 - 4ab + 6b^2$$

$$w = 6a^2 - 4ab + 4b^2$$

Diophantine equations

$$x^3 + y^3 + z^3 = w^3$$

$$(x, y, z, w) = (3, 4, 5, 6)$$

$$3^3 + 4^3 + 5^3 = 27 + 64 + 125 = 216 = 6^3$$

Parametric solution :

$$x = 3a^2 + 5ab - 5b^2$$

$$z = 5a^2 - 5ab - 3b^2$$

$$y = 4a^2 - 4ab + 6b^2$$

$$w = 6a^2 - 4ab + 4b^2$$

Diophantine equations

$$x^3 + y^3 + z^3 = w^3$$

$$(x, y, z, w) = (3, 4, 5, 6)$$

$$3^3 + 4^3 + 5^3 = 27 + 64 + 125 = 216 = 6^3$$

Parametric solution :

$$x = 3a^2 + 5ab - 5b^2$$

$$z = 5a^2 - 5ab - 3b^2$$

$$y = 4a^2 - 4ab + 6b^2$$

$$w = 6a^2 - 4ab + 4b^2$$

Ramanujan – Nagell Equation

Trygve Nagell (1895 – 1988)

$$x^2 + 7 = 2^n$$

$$1^2 + 7 = 2^3 = 8$$

$$3^2 + 7 = 2^4 = 16$$

$$5^2 + 7 = 2^5 = 32$$

$$11^2 + 7 = 2^7 = 128$$

$$181^2 + 7 = 2^{15} = 32\,768$$

Ramanujan – Nagell Equation

Trygve Nagell (1895 – 1988)

$$x^2 + 7 = 2^n$$

$$1^2 + 7 = 2^3 = 8$$

$$3^2 + 7 = 2^4 = 16$$

$$5^2 + 7 = 2^5 = 32$$

$$11^2 + 7 = 2^7 = 128$$

$$181^2 + 7 = 2^{15} = 32\,768$$

$$x^2 + D = 2^n$$

Nagell (1948) : for $D = 7$, no further solution

R. Apéry (1960) : for $D > 0$, $D \neq 7$, the equation $x^2 + D = 2^n$ has at most 2 solutions.

Examples with 2 solutions :

$$D = 23 : \quad 3^2 + 23 = 32, \quad 45^2 + 23 = 2^{11} = 2048$$

$$D = 2^{\ell+1} - 1, \ell \geq 3 : \quad (2^\ell - 1)^2 + 2^{\ell+1} - 1 = 2^{2\ell}$$

$$x^2 + D = 2^n$$

Nagell (1948) : for $D = 7$, no further solution

R. Apéry (1960) : for $D > 0$, $D \neq 7$, the equation $x^2 + D = 2^n$ has at most 2 solutions.

Examples with 2 solutions :

$$D = 23 : \quad 3^2 + 23 = 32, \quad 45^2 + 23 = 2^{11} = 2048$$

$$D = 2^{\ell+1} - 1, \ell \geq 3 : \quad (2^\ell - 1)^2 + 2^{\ell+1} - 1 = 2^{2\ell}$$

$$x^2 + D = 2^n$$

Nagell (1948) : for $D = 7$, no further solution

R. Apéry (1960) : for $D > 0$, $D \neq 7$, the equation $x^2 + D = 2^n$ has at most 2 solutions.

Examples with 2 solutions :

$$D = 23 : \quad 3^2 + 23 = 32, \quad 45^2 + 23 = 2^{11} = 2048$$

$$D = 2^{\ell+1} - 1, \ell \geq 3 : \quad (2^\ell - 1)^2 + 2^{\ell+1} - 1 = 2^{2\ell}$$

$$x^2 + D = 2^n$$

F. Beukers (1980) : at most one solution otherwise.



M. Bennett (1995) : considers the case $D < 0$.

Partitions

$$\begin{array}{ll} 1 & p(1) = 1 \\ 2 = 1 + 1 & p(2) = 2 \\ 3 = 2 + 1 = 1 + 1 + 1 & p(3) = 3 \\ 4 = 3 + 1 = 2 + 2 = 2 + 1 + 1 & \\ & = 1 + 1 + 1 + 1 & p(4) = 5 \end{array}$$

$$p(5) = 7, \quad p(6) = 11, \quad p(7) = 15, \dots$$

MacMahon : table of the first 200 values

Neil J. A. Sloane's encyclopaedia

<http://www.research.att.com/~njas/sequences/A000041>

Partitions

$$\begin{array}{ll} 1 & p(1) = 1 \\ 2 = 1 + 1 & p(2) = 2 \\ 3 = 2 + 1 = 1 + 1 + 1 & p(3) = 3 \\ 4 = 3 + 1 = 2 + 2 = 2 + 1 + 1 & \\ = 1 + 1 + 1 + 1 & p(4) = 5 \end{array}$$

$$p(5) = 7, \quad p(6) = 11, \quad p(7) = 15, \dots$$

MacMahon : table of the first 200 values

Neil J. A. Sloane's encyclopaedia

<http://www.research.att.com/~njas/sequences/A000041>

Partitions

$$\begin{array}{ll} 1 & p(1) = 1 \\ 2 = 1 + 1 & p(2) = 2 \\ 3 = 2 + 1 = 1 + 1 + 1 & p(3) = 3 \\ 4 = 3 + 1 = 2 + 2 = 2 + 1 + 1 & \\ & = 1 + 1 + 1 + 1 & p(4) = 5 \end{array}$$

$$p(5) = 7, \quad p(6) = 11, \quad p(7) = 15, \dots$$

MacMahon : table of the first 200 values

Neil J. A. Sloane's encyclopaedia

<http://www.research.att.com/~njas/sequences/A000041>

Ramanujan

$p(5n + 4)$ is a multiple of 5

$p(7n + 5)$ is a multiple of 7

$p(11n + 6)$ is a multiple of 11

$p(25n + 24)$ is a multiple of 25

$p(49n + 47)$ is a multiple of 49

$p(121n + 116)$ is a multiple of 121

Ramanujan

$p(5n + 4)$ is a multiple of 5

$p(7n + 5)$ is a multiple of 7


$p(11n + 6)$ is a multiple of 11

$p(25n + 24)$ is a multiple of 25

$p(49n + 47)$ is a multiple of 49

$p(121n + 116)$ is a multiple of 121

Partitions - Ken Ono



Manjul Bhargava, left, and Ken Ono in front of Ramanujan's house.


Honoring a Gift from Kumbakonam

Ken Ono

Today was an absolutely glorious day in Madison, Wisconsin. It is Christmas 2005, and everyone in the house is asleep after a long day of enjoying family, opening presents, and eating enormous portions of mashed potatoes and yule log cake. Yet powerful images keep me awake.

Thirty-six hours ago I returned from a six-day whirlwind journey to a far-off place. I spent forty hours on airplanes, and I endured fourteen hours in cars dodging bicycles, rickshaws, cows, goats, and masses of people on roads severely damaged by recent flooding. These floods would be blamed for at least forty-two deaths. Despite these hardships and bad luck, this adventure exceeded my lofty expectations.

I ostensibly travelled to Kumbakonam with the purpose of giving a



Bust of Srinivasa Ramanujan by artist Paul Granlund.

This adventure was a pilgrimage to pay homage to Srinivasa Ramanujan, the Indian legend whose congruences, formulas, and identities have inspired much of my own work. This fulfilled a personal journey, one with an unlikely beginning in 1984.

The Story of Ramanujan

Ramanujan was born on December 22, 1887, in Erode, a small town about 250 miles southwest of Chennai (formerly known as Madras). He was a Brahmin, a member of India's priestly caste, and as a consequence he lived his life as a strict vegetarian.

When Ramanujan was one year old, he moved to Kumbakonam, a small town about 170 miles south of Chennai, where his father Srinivasa was a cloth merchant's clerk. Kumbakonam, which is situated on the banks of the sacred Kaveri River,

Notices of the AMS, 53 N°6 (July 2006), 640-651

<http://www.ams.org/notices/200606/fea-ono.pdf>

Leonhard Euler (1707 – 1783)



$$1 + p(1)x + p(2)x^2 + \cdots + p(n)x^n + \cdots$$
$$= \frac{1}{(1-x)(1-x^2)(1-x^3)\cdots(1-x^n)\cdots}$$

$$1 + \sum_{n=1}^{\infty} p(n)x^n = \prod_{n=1}^{\infty} (1-x^n)^{-1}$$

Eulerian products

Riemann zeta function

For $s > 1$,

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_p (1 - p^{-s})^{-1}$$



Georg Friedrich Bernhard
Riemann (1826 - 1866)

Ramanujan tau function

$$x(1-x)^{-1} = \sum_{n=1}^{\infty} x^n$$

$$x \prod_{n=1}^{\infty} (1-x^n)^{24} = \sum_{n=1}^{\infty} \tau(n)x^n.$$

$$\sum_{n=1}^{\infty} \frac{\tau(n)}{n^s} = \prod_p (1 - \tau(p)p^{-s} + p^{11-2s})^{-1}$$

Ramanujan tau function

$$x(1-x)^{-1} = \sum_{n=1}^{\infty} x^n$$

$$x \prod_{n=1}^{\infty} (1-x^n)^{24} = \sum_{n=1}^{\infty} \tau(n)x^n.$$

$$\sum_{n=1}^{\infty} \frac{\tau(n)}{n^s} = \prod_p (1 - \tau(p)p^{-s} + p^{11-2s})^{-1}$$

Ramanujan tau function

$$x(1-x)^{-1} = \sum_{n=1}^{\infty} x^n$$

$$x \prod_{n=1}^{\infty} (1-x^n)^{24} = \sum_{n=1}^{\infty} \tau(n)x^n.$$

$$\sum_{n=1}^{\infty} \frac{\tau(n)}{n^s} = \prod_p (1 - \tau(p)p^{-s} + p^{11-2s})^{-1}$$

Ramanujan's Congruences

$\tau(pn)$ is divisible by p for $p = 2, 3, 5, 7, 23$.

also : congruences modulo 691
(numerator of Bernoulli number B_{12})

Ramanujan's Congruences

$\tau(pn)$ is divisible by p for $p = 2, 3, 5, 7, 23$.

also : congruences modulo 691
(numerator of Bernoulli number B_{12})

Pierre Deligne

Ramanujan's Conjecture,
proved by Deligne in 1974

$$|\tau(p)| < 2p^{11/2}$$



Hardy–Ramanujan

For *almost all* integers n , the number of prime factors of n is $\log \log n$.

$$A_\epsilon(x) = \{n \leq x ; (1 - \epsilon) \log \log n < \omega(n) < (1 + \epsilon) \log \log n\}.$$

$$\frac{1}{x} A_\epsilon(x) \rightarrow 1 \quad \text{when } x \rightarrow \infty.$$

Hardy–Ramanujan

For *almost all* integers n , the number of prime factors of n is $\log \log n$.

$$A_\epsilon(x) = \{n \leq x ; (1 - \epsilon) \log \log n < \omega(n) < (1 + \epsilon) \log \log n\}.$$

$$\frac{1}{x} A_\epsilon(x) \rightarrow 1 \quad \text{when } x \rightarrow \infty.$$

Hardy–Ramanujan

For *almost all* integers n , the number of prime factors of n is $\log \log n$.

$$A_\epsilon(x) = \{n \leq x ; (1 - \epsilon) \log \log n < \omega(n) < (1 + \epsilon) \log \log n\}.$$

$$\frac{1}{x} A_\epsilon(x) \rightarrow 1 \quad \text{when} \quad x \rightarrow \infty.$$

Highly composite numbers

(*Proc. London Math. Soc. 1915*)

$$\begin{array}{rcccccccc} n = & 2 & 4 & 6 & 12 & 24 & 36 & 48 & 60 & 120 \dots \\ d(n) = & 2 & 3 & 4 & 6 & 8 & 9 & 10 & 12 & 16 \dots \end{array}$$

Question : For $t \in \mathbf{R}$, do the conditions $2^t \in \mathbf{Z}$ and $3^t \in \mathbf{Z}$ imply $t \in \mathbf{Z}$?

Example : For $t = \frac{\log 1729}{\log 2}$, we have $2^t = 1729 \in \mathbf{Z}$, but

$$3^t = \exp((\log 3)(\log 1729)/\log 2) = 135451.447153 \dots$$

is not an integer.

Highly composite numbers

(Proc. London Math. Soc. 1915)

$$\begin{array}{rcccccccc} n = & 2 & 4 & 6 & 12 & 24 & 36 & 48 & 60 & 120 \dots \\ d(n) = & 2 & 3 & 4 & 6 & 8 & 9 & 10 & 12 & 16 \dots \end{array}$$

Question : For $t \in \mathbf{R}$, do the conditions $2^t \in \mathbf{Z}$ and $3^t \in \mathbf{Z}$ imply $t \in \mathbf{Z}$?

Example : For $t = \frac{\log 1729}{\log 2}$, we have $2^t = 1729 \in \mathbf{Z}$, but

$$3^t = \exp((\log 3)(\log 1729)/\log 2) = 135451.447153 \dots$$

is not an integer.

Highly composite numbers

(Proc. London Math. Soc. 1915)

$$\begin{array}{rcccccccc} n = & 2 & 4 & 6 & 12 & 24 & 36 & 48 & 60 & 120 \dots \\ d(n) = & 2 & 3 & 4 & 6 & 8 & 9 & 10 & 12 & 16 \dots \end{array}$$

Question : For $t \in \mathbf{R}$, do the conditions $2^t \in \mathbf{Z}$ and $3^t \in \mathbf{Z}$ imply $t \in \mathbf{Z}$?

Example : For $t = \frac{\log 1729}{\log 2}$, we have $2^t = 1729 \in \mathbf{Z}$, but

$$3^t = \exp((\log 3)(\log 1729)/\log 2) = 135451.447153 \dots$$

is not an integer.

Pàl Erdős



Carl Ludwig Siegel



Alaoglu and Erdős : *On highly composite and similar numbers*, 1944.

C.L. Siegel : For $t \in \mathbf{R}$, the conditions $2^t \in \mathbf{Z}$, $3^t \in \mathbf{Z}$ and $5^t \in \mathbf{Z}$ imply $t \in \mathbf{Z}$.

Serge Lang, K. Ramachandra : *six exponentials theorem, four exponentials conjecture*.

Five exponentials Theorem and generalizations

1985 : Five exponentials
Theorem.

1993, [Damien Roy](#), Matrices
whose coefficients are linear
forms in logarithms.
J. Number Theory **41** (1992),
no. 1, 22–47.



Approximation for π due to Ramanujan

$$\frac{63}{25} \left(\frac{17 + 15\sqrt{5}}{7 + 15\sqrt{5}} \right) = 3.141592653\ 80568820189839000630\dots$$

$$\pi = 3.141592653\ 58979323846264338328\dots$$

Another formula due to Ramanujan for π

$$\pi = \frac{9801}{\sqrt{8}} \left(\sum_{n=0}^{\infty} \frac{(4n)!(1103 + 26390n)}{(n!)^4 396^{4n}} \right)^{-1}$$

$n = 0$: 6 exact digits for 3.141592...

$n \rightarrow n + 1$: 8 more digits

Ramanujan's formula for $1/\pi$

$$\frac{1}{\pi} = \sum_{m=0}^{\infty} \binom{2m}{m} \frac{42m + 5}{2^{12m+4}}$$

Another formula due to Ramanujan for π

$$\pi = \frac{9801}{\sqrt{8}} \left(\sum_{n=0}^{\infty} \frac{(4n)!(1103 + 26390n)}{(n!)^4 396^{4n}} \right)^{-1}$$

$n = 0$: 6 exact digits for 3.141592...

$n \rightarrow n + 1$: 8 more digits

Ramanujan's formula for $1/\pi$

$$\frac{1}{\pi} = \sum_{m=0}^{\infty} \binom{2m}{m} \frac{42m + 5}{2^{12m+4}}$$

Another formula due to Ramanujan for π

$$\pi = \frac{9801}{\sqrt{8}} \left(\sum_{n=0}^{\infty} \frac{(4n)!(1103 + 26390n)}{(n!)^4 396^{4n}} \right)^{-1}$$

$n = 0$: 6 exact digits for 3.141592...

$n \rightarrow n + 1$: 8 more digits

Ramanujan's formula for $1/\pi$

$$\frac{1}{\pi} = \sum_{m=0}^{\infty} \binom{2m}{m} \frac{42m + 5}{2^{12m+4}}$$

Another formula due to Ramanujan for π

$$\pi = \frac{9801}{\sqrt{8}} \left(\sum_{n=0}^{\infty} \frac{(4n)!(1103 + 26390n)}{(n!)^4 396^{4n}} \right)^{-1}$$

$n = 0$: 6 exact digits for 3.141592...

$n \rightarrow n + 1$: 8 more digits

Ramanujan's formula for $1/\pi$

$$\frac{1}{\pi} = \sum_{m=0}^{\infty} \binom{2m}{m} \frac{42m + 5}{2^{12m+4}}$$

Decimals of π

Ramanujan's formulae were used in 1985 : $1.7 \cdot 10^7$ digits for π (1.7 crores)

In 1999 : $2 \cdot 10^{10}$ digits (2 000 crores)

18 Aug 2009 : Pi Calculation Record Destroyed : 2.5 Trillion Decimals ($2.5 \cdot 10^{12}$).

2,576,980,377,524 decimal places in 73 hours 36 minutes

Massive parallel computer called : T2K Tsukuba System.

Team leader professor Daisuke Takahashi.

Decimals of π

Ramanujan's formulae were used in 1985 : $1.7 \cdot 10^7$ digits for π (1.7 crores)

In 1999 : $2 \cdot 10^{10}$ digits (2 000 crores)

18 Aug 2009 : Pi Calculation Record Destroyed : 2.5 Trillion Decimals ($2.5 \cdot 10^{12}$).

2,576,980,377,524 decimal places in 73 hours 36 minutes

Massive parallel computer called : T2K Tsukuba System.

Team leader professor Daisuke Takahashi.

Decimals of π

Ramanujan's formulae were used in 1985 : $1.7 \cdot 10^7$ digits for π (1.7 crores)

In 1999 : $2 \cdot 10^{10}$ digits (2 000 crores)

18 Aug 2009 : Pi Calculation Record Destroyed : 2.5 Trillion Decimals ($2.5 \cdot 10^{12}$).

2,576,980,377,524 decimal places in 73 hours 36 minutes

Massive parallel computer called : T2K Tsukuba System.

Team leader professor Daisuke Takahashi.

Decimals of π

Ramanujan's formulae were used in 1985 : $1.7 \cdot 10^7$ digits for π (1.7 crores)

In 1999 : $2 \cdot 10^{10}$ digits (2 000 crores)

18 Aug 2009 : Pi Calculation Record Destroyed : 2.5 Trillion Decimals ($2.5 \cdot 10^{12}$).

2,576,980,377,524 decimal places in 73 hours 36 minutes

Massive parallel computer called : T2K Tsukuba System.

Team leader professor Daisuke Takahashi.

Ramanujan Notebooks

Written from 1903 to 1914

First : 16 chapters, 134 pages

Second : 21 chapters, 252 pages

Third : 33 pages

B.M. Wilson, G.N. Watson

Edited in 1957 in Bombay

Ramanujan Notebooks

Written from 1903 to 1914

First : 16 chapters, 134 pages

Second : 21 chapters, 252 pages

Third : 33 pages

B.M. Wilson, G.N. Watson

Edited in 1957 in Bombay

The lost notebook


George Andrews, 1976



Bruce Berndt, 1985–87 (5 volumes)

Last work of Ramanujan

Mock theta functions

-  S. ZWEGERS – « Mock ϑ -functions and real analytic modular forms. », in *Berndt, Bruce C. (ed.) et al., q-series with applications to combinatorics, number theory, and physics. Proceedings of a conference, University of Illinois, Urbana-Champaign, IL, USA, October 26-28, 2000. Providence, RI : American Mathematical Society (AMS). Contemp. Math. 291, 269-277, 2001.*

SASTRA Ramanujan Prize



SASTRA Ramanujan Prize
2009 : Kathrin Bringmann.


International Conference in
Number Theory & Mock
Theta Function
Srinivasa Ramanujan Center,
Sastra University,
Kumbakonam, Dec. 22, 2009.

Previous SASTRA Ramanujan Prize winners

Manjul Bhargava and Kannan Soundararajan (2005)



Terence Tao (2006) and Ben Green (2007)



Akshay Venkatesh (2008)



Sastra Ramanujan Prize

Year	Name	University
2005	Manjul Bhargava	Princeton University
	Kannan Soundararajan	University of Michigan
2006	Terence Tao	University of California at Los Angeles
2007	Ben Green	Cambridge University
2008	Akshay Venkatesh	Stanford University
2009	Kathrin Bringmann	University of Cologne,
		University of Minnesota
2010	Wei Zhang	Harvard University
2011	Roman Holowinsky ^[1]	Ohio State University
2012	Zhiwei Yun ^[2]	Stanford University
2013	Peter Scholze ^[3]	University of Bonn
2014	James Maynard ^[4]	Oxford University, England, and
		University of Montreal, Canada
2015	Jacob Tsimerman ^[5]	University of Toronto, Canada

https://en.wikipedia.org/wiki/SASTRA_Ramanujan_Prize

ICTP Ramanujan Prize

- 2005 **Marcelo Viana**, Brazil^[3]
- 2006 **Ramdorai Sujatha**, India^[4]
- 2007 **Jorge Lauret**, Argentina^[5]
- 2008 **Enrique Pujals**, Argentina/Brazil^[6]
- 2009 **Ernesto Lupercio**, Mexico^[7]
- 2010 **Shi Yuguang**, China^[8]
- 2011 **Philibert Nang**, Gabon^[9]
- 2012 **Fernando Codá Marques**, Brazil^[10]
- 2013 **Tian Ye**, China^[11]
- 2014 **Miguel Walsh**, Argentina^[12]
- 2015 **Amalendu Krishna**, India^[13]

https://en.wikipedia.org/wiki/ICTP_Ramanujan_Prize

ICTP Ramanujan Prize

Call for Nominations, 2016 Prize

Nomination deadline: 1 March 2016



Call for Nominations, 2016 Prize

The Ramanujan Prize for young mathematicians from developing countries has been awarded annually since 2005. The Prize is now funded by the Department of Science and Technology of the Government of India (DST), and will be administered jointly by ICTP, the International Mathematical Union (IMU), and the DST.

The Prize winner must be less than 45 years of age on 31 December of the year of the award, and have conducted outstanding research in a developing country. Researchers working in any branch of the mathematical sciences are eligible. The Prize carries a \$15,000 cash award. The winner will be invited to ICTP to receive the Prize and deliver a lecture. The Prize is usually awarded to one person, but may be shared equally among recipients who have contributed to the same body of work.

The Selection Committee will take into account not only the scientific quality of the research, but also the background of the candidate and the environment in which the work was carried out.

<https://www.ictp.it/about-ictp/prizes-awards/the-ramanujan-prize/call-for-nominations.aspx>

References (continued)

[2] Don Zagier (March 16, 2005, BNF/SMF) :

"Ramanujan to Hardy, from the first to the last letter..."

<http://smf.emath.fr/VieSociete/Rencontres/BNF/2005/>

[3] MacTutor History of Mathematics

<http://www-groups.dcs.st-and.ac.uk/~history/>

[4] Eric Weisstein worlds of mathematics, Wolfram Research

<http://scienceworld.wolfram.com/>

[5] Wikipedia, the free encyclopedia.

<http://en.wikipedia.org/wiki/Ramanujan>

Ramanujan according to Wikipedia



Erode December 22, 1887 —
Chetput, (Madras), April 26,
1920

Landau–Ramanujan constant

Mock theta functions

Ramanujan prime

Ramanujan–Soldner constant

Ramanujan theta function

Ramanujan's sum

Rogers–Ramanujan identities

http://en.wikipedia.org/wiki/Srinivasa_Ramanujan

Landau–Ramanujan constant

In mathematics, the **Landau–Ramanujan** constant occurs in a number theory result stating that the number of positive integers less than x which are the sum of two square numbers, for large x , varies as

$$x/\sqrt{\ln(x)}.$$

The constant of proportionality is the **Landau–Ramanujan** constant, which was discovered independently by **Edmund Landau** and **Srinivasa Ramanujan**.

More formally, if $N(x)$ is the number of positive integers less than x which are the sum of two squares, then

$$\lim_{x \rightarrow \infty} \frac{N(x)\sqrt{\ln(x)}}{x} \approx 0.76422365358922066299069873125.$$

Ramanujan primes

In mathematics, a **Ramanujan prime** is a prime number that satisfies a result proven by **Srinivasa Ramanujan** relating to the prime-counting function.

$$\pi(x) - \pi(x/2) \geq 1, 2, 3, 4, 5, \dots \quad \text{for all } x \geq 2, 11, 17, 29, 41, \dots$$

respectively, where $\pi(x)$ is the prime-counting function, that is, the number of primes less than or equal to x .

Ramanujan primes

In mathematics, a **Ramanujan prime** is a prime number that satisfies a result proven by **Srinivasa Ramanujan** relating to the prime-counting function.

$$\pi(x) - \pi(x/2) \geq 1, 2, 3, 4, 5, \dots \quad \text{for all } x \geq 2, 11, 17, 29, 41, \dots$$

respectively, where $\pi(x)$ is the prime-counting function, that is, the number of primes less than or equal to x .

Ramanujan primes

2, 11, 17, 29, 41, 47, 59, 67, 71, 97, 101, 107, 127, 149,
151, 167, 179, 181, 227, 229, 233, 239, 241, 263, 269, 281,
307, 311, 347, 349, 367, 373, 401, 409, 419, 431, 433, 439,
461, 487, 491, 503, 569, 571, 587, 593, 599, 601, 607, 641, ...

$a(n)$ is the smallest number such that if $x \geq a(n)$, then $\pi(x) - \pi(x/2) \geq n$, where $\pi(x)$ is the number of primes $\leq x$.

Neil J. A. Sloane's encyclopaedia

http:

[//www.research.att.com/~njas/sequences/A104272](http://www.research.att.com/~njas/sequences/A104272)

Ramanujan primes

2, 11, 17, 29, 41, 47, 59, 67, 71, 97, 101, 107, 127, 149,
151, 167, 179, 181, 227, 229, 233, 239, 241, 263, 269, 281,
307, 311, 347, 349, 367, 373, 401, 409, 419, 431, 433, 439,
461, 487, 491, 503, 569, 571, 587, 593, 599, 601, 607, 641, ...

$a(n)$ is the smallest number such that if $x \geq a(n)$, then $\pi(x) - \pi(x/2) \geq n$, where $\pi(x)$ is the number of primes $\leq x$.

Neil J. A. Sloane's encyclopaedia

http:

[//www.research.att.com/~njas/sequences/A104272](http://www.research.att.com/~njas/sequences/A104272)

Ramanujan primes

2, 11, 17, 29, 41, 47, 59, 67, 71, 97, 101, 107, 127, 149,
151, 167, 179, 181, 227, 229, 233, 239, 241, 263, 269, 281,
307, 311, 347, 349, 367, 373, 401, 409, 419, 431, 433, 439,
461, 487, 491, 503, 569, 571, 587, 593, 599, 601, 607, 641, ...

$a(n)$ is the smallest number such that if $x \geq a(n)$, then $\pi(x) - \pi(x/2) \geq n$, where $\pi(x)$ is the number of primes $\leq x$.

Neil J. A. Sloane's encyclopaedia

http:

[//www.research.att.com/~njas/sequences/A104272](http://www.research.att.com/~njas/sequences/A104272)

Ramanujan–Soldner constant



In mathematics, the Ramanujan–Soldner constant is a mathematical constant defined as the unique positive zero of the logarithmic integral function. It is named after Srinivasa Ramanujan and Johann Georg von Soldner (16 July 1776 - 13 May 1833).

Its value is approximately

1.451369234883381050283968485892027449493...

Ramanujan's sum (1918)

In number theory, a branch of mathematics, Ramanujan's sum, usually denoted $c_q(n)$, is a function of two positive integer variables q and n defined by the formula

$$c_q(n) = \sum_{\substack{1 \leq a \leq q \\ (a,q)=1}} e^{2\pi i \frac{a}{q} n}.$$

Ramanujan's sums are used in the proof of Vinogradov's theorem that every sufficiently-large odd number is the sum of three primes.

Ramanujan theta function

$$f(a, b) = \sum_{n=-\infty}^{\infty} a^{n(n+1)/2} b^{n(n-1)/2}$$



Carl Gustav Jacob Jacobi
(1804–1851)

In mathematics, the **Ramanujan** theta function generalizes the form of the **Jacobi** theta functions, while capturing their general properties. In particular, the **Jacobi** triple product takes on a particularly elegant form when written in terms of the **Ramanujan** theta.

Rogers–Ramanujan identities



Leonard James Rogers
1862 - 1933

In mathematics, the **Rogers–Ramanujan** identities are a set of identities related to basic hypergeometric series. They were discovered by **Leonard James Rogers** (1894) and subsequently rediscovered by **Srinivasa Ramanujan** (1913) as well as by **Issai Schur** (1917).

G.H. Hardy : Divergent Series



In

$$\frac{1}{1-z} = 1 + z + z^2 + z^3 + \dots$$

set $z = -1$, as Euler does :

$$1 - 1 + 1 - 1 + \dots = \frac{1}{2} \dots$$

Similarly, from the derivative of the previous series

$$\frac{1}{(1-z)^2} = 1 + 2z + 3z^2 + 4z^3 + \dots$$

deduce

$$1 - 2 + 3 - 4 + \dots = \frac{1}{4}$$

G.H. Hardy : Divergent Series



In

$$\frac{1}{1-z} = 1 + z + z^2 + z^3 + \dots$$

set $z = -1$, as Euler does :

$$1 - 1 + 1 - 1 + \dots = \frac{1}{2} \dots$$

Similarly, from the derivative of the previous series

$$\frac{1}{(1-z)^2} = 1 + 2z + 3z^2 + 4z^3 + \dots$$

deduce

$$1 - 2 + 3 - 4 + \dots = \frac{1}{4}$$

$$s = 1 - 2 + 3 - 4 + \dots$$

There are further reasons to attribute the value $1/4$ to s . For instance

$$s = 1 - (1 - 1 + 1 - 1 + \dots) - (1 - 2 + 3 - 4 + \dots) = 1 - \frac{1}{2} - s$$

gives $2s = 1/2$, hence $s = 1/4$.

Also computing the square by expanding the product

$$(1 - 1 + 1 - 1 + \dots)^2 = (1 - 1 + 1 - 1 + \dots)(1 - 1 + 1 - 1 + \dots)$$

yields to

$$1 - 2 + 3 - 4 + \dots = \left(\frac{1}{2}\right)^2 = \frac{1}{4}.$$

$$s = 1 - 2 + 3 - 4 + \dots$$

There are further reasons to attribute the value $1/4$ to s . For instance

$$s = 1 - (1 - 1 + 1 - 1 + \dots) - (1 - 2 + 3 - 4 + \dots) = 1 - \frac{1}{2} - s$$

gives $2s = 1/2$, hence $s = 1/4$.

Also computing the square by expanding the product

$$(1 - 1 + 1 - 1 + \dots)^2 = (1 - 1 + 1 - 1 + \dots)(1 - 1 + 1 - 1 + \dots)$$

yields to

$$1 - 2 + 3 - 4 + \dots = \left(\frac{1}{2}\right)^2 = \frac{1}{4}.$$

Cesaro convergence

For a series

$$a_0 + a_1 + \cdots + a_n + \cdots = s$$

converging (in the sense of **Cauchy**), the partial sums

$$s_n = a_0 + a_1 + \cdots + a_n$$

have a mean value

$$\frac{s_0 + \cdots + s_n}{n + 1}$$

which is a sequence which converges (in the sense of **Cauchy**) to s .

For the diverging series

$$1 - 1 + 1 - 1 + \cdots$$

the limit exists and is $1/2$.

Cesaro convergence

For a series

$$a_0 + a_1 + \cdots + a_n + \cdots = s$$

converging (in the sense of **Cauchy**), the partial sums

$$s_n = a_0 + a_1 + \cdots + a_n$$

have a mean value

$$\frac{s_0 + \cdots + s_n}{n + 1}$$

which is a sequence which converges (in the sense of **Cauchy**) to s .

For the diverging series

$$1 - 1 + 1 - 1 + \cdots$$

the limit exists and is $1/2$.

Cesaro convergence

Ernesto Cesàro
(1859 – 1906)



For the series

$$1 + 0 - 1 + 1 + 0 - 1 + \dots$$

the Cesaro limit

$$\lim_{n \rightarrow \infty} \frac{s_0 + \dots + s_n}{n + 1}$$

exists and is $2/3$.

Rules for summing divergent series

$$a_0 + a_1 + a_2 + \cdots = s \quad \text{implies} \quad ka_0 + ka_1 + ka_2 + \cdots = ks.$$

$$a_0 + a_1 + a_2 + \cdots = s \quad \text{and} \quad b_0 + b_1 + b_2 + \cdots = t \quad \text{implies} \\ a_0 + b_0 + a_1 + b_1 + a_2 + b_2 + \cdots = s + t.$$

$$a_0 + a_1 + a_2 + \cdots = s \quad \text{if and only if} \quad a_1 + a_2 + \cdots = s - a_0.$$

Rules for summing divergent series

$$a_0 + a_1 + a_2 + \cdots = s \quad \text{implies} \quad ka_0 + ka_1 + ka_2 + \cdots = ks.$$

$$a_0 + a_1 + a_2 + \cdots = s \quad \text{and} \quad b_0 + b_1 + b_2 + \cdots = t \quad \text{implies} \\ a_0 + b_0 + a_1 + b_1 + a_2 + b_2 + \cdots = s + t.$$

$$a_0 + a_1 + a_2 + \cdots = s \quad \text{if and only if} \quad a_1 + a_2 + \cdots = s - a_0.$$

Rules for summing divergent series

$$a_0 + a_1 + a_2 + \cdots = s \quad \text{implies} \quad ka_0 + ka_1 + ka_2 + \cdots = ks.$$

$$a_0 + a_1 + a_2 + \cdots = s \quad \text{and} \quad b_0 + b_1 + b_2 + \cdots = t \quad \text{implies} \\ a_0 + b_0 + a_1 + b_1 + a_2 + b_2 + \cdots = s + t.$$

$$a_0 + a_1 + a_2 + \cdots = s \quad \text{if and only if} \quad a_1 + a_2 + \cdots = s - a_0.$$

$$1^2 - 2^2 + 3^2 - 4^2 + \dots$$

Recall

$$\frac{1}{(1-z)^2} = 1 + 2z + 3z^2 + 4z^3 + \dots$$

Take one more derivative, you find also

$$1 \cdot 2 - 2 \cdot 3 + 3 \cdot 4 - 4 \cdot 5 + \dots = \frac{1}{4}$$

from which you deduce

$$1^2 - 2^2 + 3^2 - 4^2 + \dots = 0.$$

Further examples of divergent Series

$$1 + 1 + 1 + \dots = 0$$

$$1 - 2 + 4 - 8 + \dots = \frac{1}{3}$$

$$1 + 2 + 4 + 8 + \dots = -1$$

$$1^{2k} + 2^{2k} + 3^{2k} + 4^{2k} + 5^{2k} + \dots = 0 \quad \text{for } k \geq 1.$$

Euler :

$$1 - 1! + 2! - 3! + 4! + \dots = -e\left(\gamma - 1 + \frac{1}{2 \cdot 2!} - \frac{1}{3 \cdot 3!} + \dots\right)$$

gives the value 0.5963... also found by Ramanujan.

Further examples of divergent Series

$$1 + 1 + 1 + \dots = 0$$

$$1 - 2 + 4 - 8 + \dots = \frac{1}{3}$$

$$1 + 2 + 4 + 8 + \dots = -1$$

$$1^{2k} + 2^{2k} + 3^{2k} + 4^{2k} + 5^{2k} + \dots = 0 \quad \text{for } k \geq 1.$$

Euler :

$$1 - 1! + 2! - 3! + 4! + \dots = -e\left(\gamma - 1 + \frac{1}{2 \cdot 2!} - \frac{1}{3 \cdot 3!} + \dots\right)$$

gives the value $0.5963\dots$ also found by [Ramanujan](#).

Ramanujan's method (following Joseph Oesterlé)

Here is Ramanujan's method for computing the value of divergent series and for accelerating the convergence of series.

The series

$$a_0 - a_1 + a_2 - a_3 + \cdots$$

can be written

$$\frac{1}{2} (a_0 + (b_0 - b_1 + b_3 - b_4 + \cdots))$$

where $b_n = a_n - a_{n+1}$.

Acceleration of convergence

For instance in the case $a_n = 1/n^s$ we have $b_n \sim s/n^{s+1}$.
Repeating the process yields the analytic continuation of the **Riemann** zeta function.

For $s = -k$ where k is a positive integer, **Ramanujan's** method yields the **Bernoulli** numbers.

In the case of convergent series like

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots = \log 2,$$

or for **Euler** constant, **Ramanujan's** method gives an efficient way of accelerating the convergence.

Acceleration of convergence

For instance in the case $a_n = 1/n^s$ we have $b_n \sim s/n^{s+1}$.
Repeating the process yields the analytic continuation of the **Riemann** zeta function.

For $s = -k$ where k is a positive integer, **Ramanujan's** method yields the **Bernoulli** numbers.

In the case of convergent series like

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots = \log 2,$$

or for **Euler** constant, **Ramanujan's** method gives an efficient way of accelerating the convergence.

Acceleration of convergence

For instance in the case $a_n = 1/n^s$ we have $b_n \sim s/n^{s+1}$.
Repeating the process yields the analytic continuation of the **Riemann** zeta function.

For $s = -k$ where k is a positive integer, **Ramanujan's** method yields the **Bernoulli** numbers.

In the case of convergent series like

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots = \log 2,$$

or for **Euler** constant, **Ramanujan's** method gives an efficient way of accelerating the convergence.

Srinivasa Ramanujan

His life and his work

Michel Waldschmidt

Institut de Mathématiques de Jussieu & Paris VI

<http://www.imj-prg.fr/~michel.waldschmidt/>