

POINCARÉ DUALITY FOR HOPF ALGEBROIDS

SOPHIE CHEMLA, SORBONNE UNIVERSITÉ

ABSTRACT. We prove a twisted Poincaré duality for (full) Hopf algebroids with bijective antipode. As an application, we recover the Hochschild twisted Poincaré duality of Van Den Bergh ([27]). We also get a Poisson twisted Poincaré duality, which was already stated for oriented Poisson manifolds in [8].

1. INTRODUCTION

Left bialgebroids over a (possibly) non-commutative basis A generalize bialgebras. If U is a left bialgebroid, there is a natural U -module structure on A and the category of left modules over a left bialgebroid U is monoidal. Nevertheless, A is generally not a right U -module. *Left Hopf left bialgebroids* (or \times_A -Hopf algebras [24]) generalize Hopf algebras. In a left Hopf left bialgebroid U , the existence of an antipode is not required but, for any element $u \in U$, there exists an element $u_+ \otimes u_-$ corresponding to $u_{(1)} \otimes S(u_{(2)})$. The more restrictive structure of *full Hopf algebroids* ([2]) ensures the existence of an antipode. If L is a Lie-Rinehart algebra (or Lie algebroid) over a commutative k -algebra A ([23]), there exists a standard left bialgebroid structure on its enveloping algebra $V(L)$. This structure is left Hopf. Kowalzig showed ([17]) that $V(L)$ is a full Hopf algebroid if and only if there exists a right $V(L)$ -module structure on A . If X is a C^∞ Poisson manifold and $A = C^\infty(X)$, the A -module of global differential one forms $\Omega^1(X)$ is endowed with a natural Lie-Rinehart structure over A , which is of much interest ([5], [9], [11], [15], [22], [28] etc...). In particular, Huebschmann ([11]) exhibited a right $V(\Omega^1(X))$ -module structure on A (denoted A_P) that makes $V(\Omega^1(X))$ a full Hopf algebroid. He also interpreted the Lichnerowicz Poisson cohomology $H_{Pois}^i(X)$ as $\text{Ext}_{V(\Omega^1(X))}^i(A, A)$ and the Poisson homology $H_i^{Pois}(X)$ ([4], [16]) of X as $\text{Tor}_{V(\Omega^1(X))}(A_P, A)$.

A *Poincaré duality* theorem was proved in [5] for Lie-Rinehart algebras and then extended to left Hopf left bialgebroids in [18]. It asserts, under some conditions, that if $\text{Ext}_U^i(A, U) = 0$ for $i \neq d$ then, for all left U -modules M and all $n \in \mathbb{N}$, there is an isomorphism

$$\text{Ext}_U^n(A, M) \simeq \text{Tor}_{d-n}^U(M \otimes_A \Lambda, A),$$

where $\Lambda := \text{Ext}_U^d(A, U)$ is endowed with the right U -module structure given by right multiplication in U . If $U = V(L)$ is the enveloping algebra of a finitely generated

projective Lie-Rinehart algebra L , it is shown in [5] that $\text{Ext}_{V(L)}^n(A, V(L)) = 0$ if $n \neq \dim L$. Moreover, $\text{Ext}_{V(L)}^{\dim L}(A, V(L)) \simeq \Lambda_A^{\dim L}(L^*)$.

We give a new formulation of Poincaré duality in the case where U as well as its coopposite U_{coop} is left Hopf and A is endowed with a right U -module structure (denoted A_R) such that the the A^e -module $\blacktriangleright A_R \blacktriangleleft$ is invertible.

Theorem 3.5 *Let U be a left and right Hopf left bialgebroid over A . Assume the following:*

- (i) $\text{Ext}_U^i(A, U) = \{0\}$ if $i \neq d$ and set $\Lambda = \text{Ext}_U^d(A, U)$.
- (ii) The left U -module A admits a finitely generated projective resolution of finite length.
- (iii) A is endowed with a right U -module structure (denoted A_R) such that the A^e -module $\blacktriangleright A_R \blacktriangleleft$ is invertible.
- (iv) Let \mathcal{T} be the left U -module $\text{Hom}_A(A_R \blacktriangleleft, \Lambda \blacktriangleleft)$ (see Proposition 2.7). The A -module $\triangleright \mathcal{T}$ and the A^{op} -module $\mathcal{T} \triangleleft$ are projective.

Then, for all left U -modules M and all $i \in \mathbb{N}$, there is an isomorphism

$$\text{Ext}_U^i(A, M) \simeq \text{Tor}_{d-i}^U(A_R, \mathcal{T} \triangleleft \otimes_A \triangleright M).$$

Assume now that H is a full Hopf algebroid. The antipode allows us to transform any left (resp. right) H -module M (resp. N) into a right (resp. left) H -module denoted M_S (resp. ${}_S N$). Thus from the left H -module structure on A , we can construct a right H -module structure A_S . From the right H -module structure on Λ , we can make a left H -module structure denoted ${}_S \Lambda$. The duality states the following:

$$\text{Ext}_H^i(A, M) \simeq \text{Tor}_{d-i}^H(A_S, {}_S \Lambda \otimes_A M).$$

In the special case of the (full) Hopf algebroid $A \otimes A^{op}$, we recover the Hochschild twisted Poincaré duality of [27]. In the special case where X is a Poisson manifold and $H = V(\Omega^1(X))$, the duality above can be rewritten in terms of Poisson cohomology and homology. Let M be a left H -module. The coproduct on H allows us to endow ${}_S \Lambda \otimes_A M$ with a left H -module structure. Denote by $H_{Pois}^i(M)$ the Poisson cohomology with coefficients in M and let $H_i^{Pois}({}_S \Lambda \otimes_A M)$ denote the Poisson homology with coefficients in ${}_S \Lambda \otimes_A M$. There is an isomorphism

$$H_{Pois}^i(M) \simeq H_{d-i}^{Pois}({}_S \Lambda \otimes_A M).$$

This formula was stated in [8] for oriented Poisson manifolds (see also [20] for polynomial algebras with quadratic Poisson structures, [30] for linear Poisson structures, [21] for general polynomial Poisson algebras).

Notations

Fix an (associative, unital, commutative) ground ring k . Unadorned tensor products will always be meant over k . All other algebras, modules etc. will have an underlying structure of a k -module. Secondly, fix an associative and unital k -algebra A , i.e., a ring with a ring homomorphism $\eta_A : k \rightarrow Z(A)$ to its centre.

Denote by A^{op} the opposite algebra and by $A^e := A \otimes A^{\text{op}}$ the enveloping algebra of A , and by $A\text{-Mod}$ the category of left A -modules.

The notions of A -ring and A -coring are direct generalizations of the notions of algebra and coalgebra over a commutative ring.

Definition 1.1. *An A -coring is a triple (C, Δ, ϵ) where C is an A^e -module (with left action L_A and right action R_A), $\Delta : C \rightarrow C \otimes_A C$ and $\epsilon : C \rightarrow A$ are A^e -module morphisms such that*

$$(\Delta \otimes \text{id}_C) \circ \Delta = (\text{id}_C \otimes \Delta) \circ \Delta, \quad L_A \circ (\epsilon \otimes \text{id}_C) \circ \Delta = \text{id}_C = R_A \circ (\text{id}_C \otimes \epsilon) \circ \Delta.$$

As usual, we adopt Sweedler's Σ -notation $\Delta(c) = c_{(1)} \otimes c_{(2)}$ or $\Delta(c) = c^{(1)} \otimes c^{(2)}$ for $c \in C$.

The notion of A -ring is dual to that of A -coring. It is well known (see [1]) that A -rings H correspond bijectively to k -algebra homomorphisms $\iota : A \rightarrow H$. An A -ring H is endowed with the following A^e -module structure :

$$\forall h \in H, \quad a, b \in H, \quad a \cdot h \cdot b = \iota(a)h\iota(b).$$

2. PRELIMINARIES

We recall the notions and results with respect to bialgebroids that are needed to make this article self content; see, *e.g.*, [17] and references below for an overview on this subject.

2.1. Bialgebroids. For an A^e -ring U given by the k -algebra map $\eta : A^e \rightarrow U$, consider the restrictions $s := \eta(- \otimes 1_U)$ and $t := \eta(1_U \otimes -)$, called *source* and *target* map, respectively. Thus an A^e -ring U carries two A -module structures from the left and two from the right, namely

$$a \triangleright u \triangleleft b := s(a)t(b)u, \quad a \blacktriangleright u \blacktriangleleft b := ut(a)s(b), \quad \forall a, b \in A, u \in U.$$

If we let $U_{\triangleleft} \otimes_A \triangleright U$ be the corresponding tensor product of U (as an A^e -module) with itself, we define the (*left*) *Takeuchi-Sweedler product* as

$$U_{\triangleleft} \times_A \triangleright U := \left\{ \sum_i u_i \otimes u'_i \in U_{\triangleleft} \otimes_A \triangleright U \mid \sum_i (a \blacktriangleright u_i) \otimes u'_i = \sum_i u_i \otimes (u'_i \blacktriangleleft a), \forall a \in A \right\}.$$

By construction, $U_{\triangleleft} \times_A \triangleright U$ is an A^e -submodule of $U_{\triangleleft} \otimes_A \triangleright U$; it is also an A^e -ring via factorwise multiplication, with unit $1_U \otimes 1_U$ and $\eta_{U_{\triangleleft} \times_A \triangleright U}(a \otimes \tilde{a}) := s(a) \otimes t(\tilde{a})$.

Symmetrically, one can consider the tensor product $U_{\blacktriangleleft} \otimes_A \blacktriangleright U$ and define the (*right*) *Takeuchi-Sweedler product* as $U_{\blacktriangleleft} \times_A \blacktriangleright U$, which is an A^e -ring inside $U_{\blacktriangleleft} \otimes_A \blacktriangleright U$.

Definition 2.1. ([26]) *A left bialgebroid (U, A) is a k -module U with the structure of an A^e -ring (U, s^ℓ, t^ℓ) and an A -coring $(U, \Delta_\ell, \epsilon)$ subject to the following compatibility relations:*

- (i) *The A^e -module structure on the A -coring U is that of $\triangleright U_{\triangleleft}$.*
- (ii) *The coproduct Δ_ℓ is a unital k -algebra morphism taking values in $U_{\triangleleft} \times_A \triangleright U$.*

(iii) For all $a, b \in A$ and $u, u' \in U$, one has:

$$\epsilon(1_U) = 1_A, \quad \epsilon(a \triangleright u \triangleleft b) = a\epsilon(u)b, \quad \epsilon(uu') = \epsilon(u \triangleleft \epsilon(u')) = \epsilon(\epsilon(u') \triangleright u). \quad (2.1)$$

A morphism between left bialgebroids (U, A) and (U', A') is a pair (F, f) of maps $F : U \rightarrow U'$, $f : A \rightarrow A'$ that commute with all structure maps in an obvious way.

As for any ring, we can define the categories $U - \text{Mod}$ and $\text{Mod} - U$ of left and right modules over U . Note that $U - \text{Mod}$ forms a monoidal category but $\text{Mod} - U$ usually does not. However, in both cases there is a forgetful functor $U - \text{Mod} \rightarrow A^e - \text{Mod}$, respectively $\text{Mod} - U \rightarrow A^e - \text{Mod}$ given by the following formulas: For $m \in M$, $n \in N$ and $a, b \in A$,

$$a \triangleright m \triangleleft b := s^\ell(a)t^\ell(b)m, \quad a \triangleright m \triangleleft b := ns^\ell(b)t^\ell(a).$$

For example, the base algebra A itself is a left U -module via the left action

$$u(a) := \epsilon(u \triangleleft a) = \epsilon(a \triangleright u), \quad \forall u \in U, \quad \forall a \in A, \quad (2.2)$$

but in general, there is no right U -action on A .

Example 2.2. Let A be a commutative k -algebra and $\text{Der}_k(A)$ the A -module of k -derivations of A . Let L be a Lie-Rinehart algebra ([23]) over A with anchor $\rho : L \rightarrow \text{Der}_k(A)$. Its enveloping algebra $V(L)$ is endowed with a standard left bialgebroid ([28]) described by the following: For all $a \in A$, $D \in L$ and $u \in V(L)$,

- (i) s^ℓ and t^ℓ are equal to the natural injection $\iota : A \rightarrow V(L)$;
- (ii) $\Delta_\ell : V(L) \rightarrow V(L) \otimes_A V(L)$, $\Delta_\ell(a) = a \otimes_A 1$, $\Delta_\ell(D) = D \otimes_A 1 + 1 \otimes_A D$;
- (iii) $\epsilon(u) = \rho(u)(1)$.

In this example, the left action of $V(L)$ on A coincides with the anchor extended to $V(L)$.

2.2. Left and right Hopf left bialgebroids. For any left bialgebroid U , define the Hopf-Galois maps

$$\begin{aligned} \alpha_\ell : \triangleright U \otimes_{A^{\text{op}}} U \triangleleft &\rightarrow U \triangleleft \otimes_A \triangleright U, & u \otimes_{A^{\text{op}}} v &\mapsto u_{(1)} \otimes_A u_{(2)} v, \\ \alpha_r : U \triangleleft \otimes^A \triangleright U &\rightarrow U \triangleleft \otimes_A \triangleright U, & u \otimes^A v &\mapsto u_{(1)} v \otimes_A u_{(2)}. \end{aligned}$$

With the help of these maps, we make the following definition due to Schauenburg [24]:

Definition 2.3. A left bialgebroid U is called a left Hopf left bialgebroid or \times_A Hopf algebra if α_ℓ is a bijection. Likewise, it is called a right Hopf left bialgebroid if α_r is a bijection. In either case, we adopt for all $u \in U$ the following (Sweedler-like) notation

$$u_+ \otimes_{A^{\text{op}}} u_- := \alpha_\ell^{-1}(u \otimes_A 1), \quad u_{[+]} \otimes^A u_{[-]} := \alpha_r^{-1}(1 \otimes_A u), \quad (2.3)$$

and call both maps $u \mapsto u_+ \otimes_{A^{\text{op}}} u_-$ and $u \mapsto u_{[+]} \otimes^A u_{[-]}$ translation maps.

Remarks 2.4. Let $(U, A, s^\ell, t^\ell, \Delta, \epsilon)$ be a left bialgebroid.

- (i) In case $A = k$ is central in U , one can show that α_ℓ is invertible if and only if U is a Hopf algebra, and the translation map reads $u_+ \otimes u_- := u_{(1)} \otimes S(u_{(2)})$, where S is the antipode of U . On the other hand, U is a Hopf algebra with invertible antipode if and only if both α_ℓ and α_r are invertible, and then $u_{[+]} \otimes u_{[-]} := u_{(2)} \otimes S^{-1}(u_{(1)})$.
- (ii) The underlying left bialgebroid in a *full* Hopf algebroid with bijective antipode is both a left and right Hopf left bialgebroid (but not necessarily vice versa); see [2] [Prop. 4.2] for the details of this construction recalled in paragraph 3.10

Example 2.5. If L is a Lie-Rinehart algebra over a commutative k -algebra A with anchor ρ , then its enveloping algebra $V(L)$, endowed with its standard bialgebroid structure, is a left Hopf left bialgebroid. The translation map is described as follows (see Proposition 2.6; in this case, $A = A^{\text{op}}$ and $s^\ell = t^\ell$) : If $a \in A$ and $D \in L$,

$$a_+ \otimes_{A^{\text{op}}} a_- = a \otimes_{A^{\text{op}}} 1, \quad D_+ \otimes_{A^{\text{op}}} D_- = D \otimes_{A^{\text{op}}} 1 - 1 \otimes_{A^{\text{op}}} D.$$

It is also a right Hopf left bialgebroid as it is cocommutative.

The following proposition collects some properties of the translation maps [24]:

Proposition 2.6. *Let U be a left bialgebroid.*

- (i) *If U is a left Hopf left bialgebroid, the following relations hold:*

$$\begin{aligned} u_+ \otimes_{A^{\text{op}}} u_- &\in U \times_{A^{\text{op}}} U, \\ u_{+(1)} \otimes_A u_{+(2)} u_- &= u \otimes_A 1 \in U_{\triangleleft} \otimes_A U_{\triangleright}, \\ u_{(1)+} \otimes_{A^{\text{op}}} u_{(1)-} u_{(2)} &= u \otimes_{A^{\text{op}}} 1 \in \blacktriangleright U \otimes_{A^{\text{op}}} U_{\triangleleft}, \\ u_{+(1)} \otimes_A u_{+(2)} \otimes_{A^{\text{op}}} u_- &= u_{(1)} \otimes_A u_{(2)+} \otimes_{A^{\text{op}}} u_{(2)-}, \\ u_+ \otimes_{A^{\text{op}}} u_{-(1)} \otimes_A u_{-(2)} &= u_{++} \otimes_{A^{\text{op}}} u_- \otimes_A u_{+-}, \\ (uv)_+ \otimes_{A^{\text{op}}} (uv)_- &= u_+ v_+ \otimes_{A^{\text{op}}} v_- u_-, \\ u_+ u_- &= s^\ell(\varepsilon(u)), \\ \varepsilon(u_-) \blacktriangleright u_+ &= u, \\ (s^\ell(a)t^\ell(b))_+ \otimes_{A^{\text{op}}} (s^\ell(a)t^\ell(b))_- &= s^\ell(a) \otimes_{A^{\text{op}}} s^\ell(b), \end{aligned}$$

where, in (i), we mean the Takeuchi-Sweedler product

$$U \times_{A^{\text{op}}} U := \{ \sum_i u_i \otimes v_i \in \blacktriangleright U \otimes_{A^{\text{op}}} U_{\triangleleft} \mid \sum_i u_i \triangleleft a \otimes v_i = \sum_i u_i \otimes a \blacktriangleright v_i, \forall a \in A \}.$$

- (ii) *There are similar relations for $u_{[+]} \otimes_A u_{[-]}$ if U is a right Hopf left bialgebroid (see [7] for an exhaustive list).*

The existence of a translation map if U is a left or right Hopf left bialgebroid makes it possible to endow Hom-spaces and tensor products of U -modules with further natural U -module structures. These structures were systematically studied in [7] (proposition 3.1.1). They generalize the case of $V(L)$ ([6], see [3], [14] for the particular case $L = \text{Der}(A)$)

Proposition 2.7. *Let (U, A) be a left bialgebroid, $M, M' \in U - \text{Mod}$ and $N, N' \in \text{Mod} - U$ be left resp. right U -modules. We denote the respective actions by juxtaposition.*

- (i) *Let (U, A) be additionally a left Hopf left bialgebroid.*

(a) The A^e -module $\text{Hom}_{A^{\text{op}}}(M, M')$ carries a left U -module structure given by

$$(u \cdot f)(m) := u_+(f(u_-m)). \quad (2.4)$$

(b) The A^e -module $\text{Hom}_A(N, N')$ carries a left U -module structure via

$$(u \cdot f)(n) := (f(nu_+))u_-. \quad (2.5)$$

(c) The A^e -module $\blacktriangleright N \otimes_{A^{\text{op}}} M \blacktriangleleft$ carries a right U -module structure via

$$(n \otimes_{A^{\text{op}}} m) \cdot u := nu_+ \otimes_{A^{\text{op}}} u_-m. \quad (2.6)$$

(ii) Let (U, A) be a right Hopf left bialgebroid instead.

(a) The A^e -module $\text{Hom}_A(M, M')$ carries a left U -module structure given by

$$(u \cdot f)(m) := u_{[+]}(f(u_{[-]}m)). \quad (2.7)$$

(b) The A^e -module $\text{Hom}_{A^{\text{op}}}(N, N')$ carries a left U -module structure given by

$$(u \cdot f)(n) := (f(nu_{[+]}))u_{[-]}. \quad (2.8)$$

(c) The A^e -module $N \blacktriangleleft \otimes^A \blacktriangleright M$ carries a right U -module structure given by

$$(n \otimes^A m) \cdot u := nu_{[+]} \otimes^A u_{[-]}m. \quad (2.9)$$

Corollary 2.8. ([7]) Let U be left and right left bialgebroid. For any $N \in \text{Mod } -U$, the evaluation map

$$P \blacktriangleleft \otimes_A \blacktriangleright \text{Hom}_A(\blacktriangleright P, \blacktriangleright N) \rightarrow N, \quad p \otimes_A \phi \mapsto \phi(p) \quad (2.10)$$

is a morphism of right U -modules.

Proof. a very similar result is stated in [7], Proposition 3.2.1. \square

3. POINCARÉ DUALITY

We start by recalling the definition of an *invertible module* ([10]).

Definition 3.1. Let A be k -algebra. An $A \otimes A^{\text{op}}$ -module X is invertible if there exists an $A \otimes A^{\text{op}}$ -module Y and isomorphisms of $A \otimes A^{\text{op}}$ -modules

$$\begin{aligned} f : X \otimes_A Y &\rightarrow A \\ g : Y \otimes_A X &\rightarrow A \end{aligned}$$

such that for all $(x, y) \in X^2$ and all $(x', y') \in Y$

$$f(x, y')y = xg(y', y) \quad \text{and} \quad x'f(x, y') = g(x', x)y'.$$

Remark 3.2. In [29], Yekutieli classifies invertible $A \otimes A^{\text{op}}$ -modules in the case where A is a non-commutative graded algebra.

Proposition 3.3. ([12] p. 167)

Let A be a k -algebra and let P be an $A \otimes A^{op}$ -module. Then, if M is an A -module, we endow $\text{Hom}_A(P, M)$ with the $A \otimes A^{op}$ -module structure: For all $(a, b) \in A$, $p \in P$ and $\lambda \in \text{Hom}_A(P, M)$,

$$\langle a \cdot \lambda \cdot b, p \rangle = \langle \lambda, p \cdot a \rangle b.$$

P an invertible A^e -module if and only if it satisfies the following conditions:

- The A -module P is finitely generated projective A -module.
- The left $A \otimes A^{op}$ -module morphism

$$g : A \rightarrow \text{Hom}_A(P, P), \quad a \mapsto \{p \mapsto p \cdot a\}$$

is an isomorphism.

- The evaluation map

$$ev : P \otimes_A \text{Hom}_A(P, A) \rightarrow A, \quad p \otimes_{A^{op}} \phi \mapsto \phi(p) \quad (3.1)$$

is an isomorphism of $A \otimes A^{op}$ -modules.

Remark 3.4. Let U be a left and right Hopf left bialgebroid over A . If moreover, P is endowed with a right U -module structure such that the $A \otimes A^{op}$ -module structure on P is isomorphic to that given by \blacktriangleright and \blacktriangleleft , then the evaluation map is an isomorphism of left U -modules (Corollary 3.1).

We can now state *twisted Poincaré duality*:

Theorem 3.5. Let U be a left and right Hopf left bialgebroid over A . Assume the following:

- (i) $\text{Ext}_U^i(A, U) = \{0\}$ if $i \neq d$ and set $\Lambda = \text{Ext}_U^d(A, U)$ with the right U -module structure given by right multiplication on U .
- (ii) The left U -module A admits a finitely generated projective resolution of finite length.
- (iii) A is endowed with a right U -module structure (denoted A_R) such that the A^e -module $\blacktriangleright A_R \blacktriangleleft$ is invertible.
- (iv) Let \mathcal{T} be the left U -module $\text{Hom}_A(\blacktriangleright A_R, \blacktriangleright \Lambda)$ (see Proposition 2.7). The A -module $\triangleright \mathcal{T}$ and the A^{op} -module $\mathcal{T} \triangleleft$ are projective.

Then, for all left U -modules M and all $n \in \mathbb{N}$, there is an isomorphism

$$\text{Ext}_U^i(A, M) \simeq \text{Tor}_{d-i}^U(A_R, \mathcal{T} \triangleleft \otimes_{A \triangleright} M).$$

Remark 3.6. In the case where U is the enveloping algebra $V(L)$ of a finitely generated projective Lie-Rinehart algebra L over A with anchor $\rho : L \rightarrow \text{Der}_k(A)$, the hypothesis are all verified (see [5]). More precisely, if L is a projective A -module with constant rank n , then $\text{Ext}_{V(L)}^i(A, V(L)) = \{0\}$ if $i \neq n$. To describe the right $V(L)$ -module $\text{Ext}_{V(L)}^n(A, V(L))$, we make use of the *Lie derivative* \mathcal{L} over the Lie Rinehart algebra L , which we briefly recall.

The k -Lie algebra L acts on $L^* = \text{Hom}_A(L, A)$ as follows : For all $D, \Delta \in L$ and $\lambda \in L^*$,

$$\mathcal{L}_D(\lambda)(\Delta) = \rho(D) [\lambda(\Delta)] - \lambda([D, \Delta]).$$

By extension, the Lie derivative \mathcal{L}_D is also defined on $\Lambda_A^n(L^*)$. This allows us to endow $\Lambda_A^n(L^*)$ with a natural right $V(L)$ -module structure determined as follows:

$$\forall a \in A, \quad \forall D \in L, \quad \forall \omega \in \Lambda_A^n(L^*), \quad \omega \cdot a = a\omega, \quad \omega \cdot D = -\mathcal{L}_D(\omega).$$

The right $V(L)$ -modules $\text{Ext}_{V(L)}^n(A, V(L))$ and $\Lambda_A^n(L^*)$ are isomorphic ([5], see [3] or [14] for the special case $L = \text{Der}_k(A)$).

In the particular case where X is a n -dimensional Poisson manifold, $A = \mathcal{C}^\infty(X)$, $L = \Omega^1(X)$, $L^* = \text{Der}(A)$, the Lie derivative \mathcal{L}_{df} over $\Lambda_A^n(L^*) = \Lambda_A^n(\text{Der}(A))$ is the Lie derivative with respect to the Hamiltonian vector field $H_f = \{f, -\}$.

Proof. To prove the theorem 3.5, we will make use of the following lemma where the U -module structures are given by Proposition 2.7:

Lemma 3.7. ([18], Lemma 16) *Let U be a right Hopf left bialgebroid. Let N be a right U -module and let M and \mathcal{T} be two left U -modules. There is an isomorphism of k -modules:*

$$(N \blacktriangleleft \otimes_{A \blacktriangleright} \mathcal{T}) \otimes_U M \simeq N \otimes_U (\mathcal{T} \blacktriangleleft \otimes_{A \blacktriangleright} M).$$

Let P^\bullet be a bounded finitely generated projective resolution of the left U -module A and let Q^\bullet be a projective resolution of the left U -module M . The following computation holds in $D^b(k - \text{Mod})$, the bounded derived category of k -modules.

$$\begin{aligned} R\text{Hom}_U(A, M) &\simeq \text{Hom}_U(P^\bullet, M) \\ &\simeq \text{Hom}_U(P^\bullet, U) \otimes_U M \\ &\simeq \Lambda[-d] \otimes_U Q^\bullet \\ &\simeq [A_R \blacktriangleleft \otimes_{A \blacktriangleright} \mathcal{T}] \otimes_U Q^\bullet[-d] \quad (\text{Remark 3.4}) \\ &\simeq A_R \otimes_U (\mathcal{T} \blacktriangleleft \otimes_{A \blacktriangleright} Q^\bullet) \quad (\text{previous lemma}) \\ &\simeq A_R \otimes_U^L (\mathcal{T} \blacktriangleleft \otimes_{A \blacktriangleright} M) \end{aligned}$$

The last isomorphism follows from the fact the A -module $\blacktriangleright \mathcal{T}$ is projective and from the lemma:

Lemma 3.8. *Denote by ${}^\ell U$ the left U -module structure on U given by left multiplication. The map*

$$\begin{aligned} \alpha_r(\mathcal{T}) : {}^\ell U \blacktriangleleft \otimes_{A \blacktriangleright} \mathcal{T} &\rightarrow \mathcal{T} \blacktriangleleft \otimes_{A \blacktriangleright} U \\ u \otimes t &\mapsto u_{(1)} t \otimes u_{(2)} \end{aligned}$$

is an isomorphism. One has $\alpha_r^{-1}(t \otimes u) = u_{[+]} \otimes u_{[-]}$. Thus the U -module $\mathcal{T} \blacktriangleleft \otimes_{A \blacktriangleright} U$ is projective if the A -module $\blacktriangleright \mathcal{T}$ is projective.

□

Remark 3.9. (i) In the case where $U = A \otimes A^{op}$ (see examples 3.12), $\text{Ext}_U^i(A, M)$ is the Hochschild cohomology and we recover Van den Berg's Hochschild twisted Poincaré duality. Moreover, the beginning of the proof is similar to that of [27] (Theorem 1).

(ii) The isomorphism $\text{Ext}_U^n(A, M) \simeq \text{Tor}_{d-n}^U(M \blacktriangleleft \otimes_{A \blacktriangleright} \Lambda, A)$ is proved in [18]. But, one can show that if the A -module $\Lambda \blacktriangleleft$ is projective, one has an isomorphism $\text{Tor}_{d-n}^U(M \blacktriangleleft \otimes_{A \blacktriangleright} \Lambda, A) \simeq \text{Tor}_{d-n}^U(\Lambda, M)$.

In the case of full Hopf algebroids, there is a natural choice of right U -module structure on A .

3.10. Reminder on full Hopf algebroids. Recall that a *full Hopf algebroid* structure ([1], [2]) on a k -module H consists of the following data:

- (i) a left bialgebroid structure $H^\ell := (H, A, s^\ell, t^\ell, \Delta_\ell, \epsilon)$ over a k -algebra A ;
- (ii) a right bialgebroid structure $H^r := (H, B, s^r, t^r, \Delta_r, \partial)$ over a k -algebra B ;
- (iii) the assumption that the k -algebra structures for H in (i) and in (ii) be the same;
- (iv) a k -module map $S : H \rightarrow H$;
- (v) some compatibility relations between the previously listed data for which we refer to *op. cit.*

The detailed definition with the same notation can be found in [19]. We shall denote by lower Sweedler indices the left coproduct Δ_ℓ and by upper indices the right coproduct Δ_r , that is, $\Delta_\ell(h) =: h_{(1)} \otimes_A h_{(2)}$ and $\Delta_r(h) =: h^{(1)} \otimes_B h^{(2)}$ for any $h \in H$. A full Hopf algebroid (with bijective antipode) is both a left and right Hopf left bialgebroid but not necessarily vice versa. In this case, the translation maps in (2.3) are given by

$$h_+ \otimes_{A^{\text{op}}} h_- = h^{(1)} \otimes_{A^{\text{op}}} S(h^{(2)}) \quad \text{and} \quad h_{[+]} \otimes_{B^{\text{op}}} h_{[-]} = h^{(2)} \otimes_{B^{\text{op}}} S^{-1}(h^{(1)}), \quad (3.2)$$

formally similar as for Hopf algebras.

The following lemma [1, 2] will be needed to prove the main result in this subsection.

Proposition 3.11. *Let $H = (H^\ell, H^r)$ be a (full) Hopf algebroid over A with bijective antipode S . Then the following statement holds:*

- (i) *The maps $\nu := \partial s^\ell : A \rightarrow B^{\text{op}}$ and $\mu := \epsilon s^r : B \rightarrow A^{\text{op}}$ are isomorphisms of k -algebras.*
- (ii) *One has $\nu^{-1} = \epsilon t^r$ and $\mu^{-1} = \partial t^\ell$.*
- (iii) *The pair of maps $(S, \nu) : H^\ell \rightarrow (H^r)_{\text{coop}}^{\text{op}}$ gives an isomorphism of left bialgebroids.*
- (iv) *The pair of maps $(S, \mu) : H^r \rightarrow (H^\ell)_{\text{coop}}^{\text{op}}$ gives an isomorphism of right bialgebroids.*

Examples 3.12. (i) Let A be a k -algebra, then $A^e = A \otimes_k A^{\text{op}}$ is a A -Hopf algebroid described as follows: For all $a, b \in A$,

- $s^\ell(a) = a \otimes_k 1, \quad t^\ell(b) = 1 \otimes_k b$;
- $\Delta_\ell : A^e \rightarrow A^e \otimes_A A^e, \quad a \otimes b \mapsto (a \otimes_k 1) \otimes_A (1 \otimes_k b)$;
- $\epsilon : A^e \rightarrow A, \quad a \otimes b \mapsto ab$;
- $s^r(a) = 1 \otimes_k a, \quad t^r(b) = b \otimes_k 1$;
- $\Delta_r : A^e \rightarrow A^e \otimes_{A^{\text{op}}} A^e, \quad a \otimes b \mapsto (1 \otimes_k a) \otimes_A (b \otimes_k 1)$;
- $\partial : A^e \rightarrow A, \quad a \otimes b \mapsto ba$.

- (ii) Let A be a commutative k -algebra and L be a Lie-Rinehart algebra over A . Its enveloping algebra $V(L)$ is endowed with a standard left bialgebroid

structure (see Example 2.2). Kowalzig ([17]) has shown that the left bialgebroid $V(L)$ can be endowed with a Hopf algebroid structure if and only if there exists a right $V(L)$ -module structure on A . Then the right bialgebroid structure $V(L)_r$ is described as follows: For any $a \in A$, $D \in L$ and $u \in V(L)$,

- (a) $\partial(u) = 1 \cdot u$;
- (b) $\Delta_r : V(L) \rightarrow V(L) \blacktriangleleft \otimes_A \blacktriangleright V(L)$, $\Delta_r(D) = D \otimes_A 1 + 1 \otimes_A D - \partial(X) \otimes_A 1$ and $\Delta_r(a) = a \otimes 1$;
- (c) $S(a) = a$, $S(D) = -D + \partial(D)$.

It is in particular the case if X is a C^∞ Poisson manifold, $A = C^\infty(X)$ and $L = \Omega^1(X)$ is the A -module of global differential 1-forms on X . Huebschmann has shown ([11]) that there is a right $V(\Omega^1(X))$ -module structure on A determined as follows: For all $(a, u, v) \in A^3$,

$$a \cdot u = au \quad \text{and} \quad a \cdot udv = \{au, v\}.$$

Thus, $V(\Omega^1(X))$ is endowed with a (full) Hopf algebroid structure.

Notation 3.13. Let (H^ℓ, H^r, S) be a full Hopf algebroid over A .

- (i) If N is a right H^ℓ -module, we will denote by ${}_S N$ the left H^ℓ -module defined by

$$\forall h \in H, \quad \forall n \in N, \quad h \cdot_S n = n \cdot S(h).$$

- (ii) If M is a left H^ℓ -module, we will denote by M_S the right H^ℓ -module defined by

$$\forall h \in H, \quad \forall m \in M, \quad m \cdot_S h = S(h) \cdot m.$$

Remark 3.14. If $H = (H^\ell, H^r, S)$ is a Hopf algebroid over a k -algebra A . We have the following module structures:

- a left H^ℓ -module structure given by $h \cdot_\ell a = \epsilon(hs^\ell(a)) = \epsilon(ht^\ell(a))$.
- a right H^r -module structure given by $\alpha \cdot_r h = \partial(s^r(\alpha)h) = \partial(t^r(a)h)$.

Thanks to the Proposition 3.11, these two structures are linked by the relation

$$S(h) \cdot_\ell \mu(\alpha) = \mu[\alpha \cdot_r h].$$

Theorem 3.15. Let (H^ℓ, H^r) be a full Hopf algebroid over A with bijective antipode S . Consider A with its left H -module structure (as in Remark 3.14). We keep the notation of Proposition 3.11, in particular $\mu = \epsilon s^r$ and $\nu = \partial s^\ell$.

- (i) If $a \in A$, then $1 \cdot_S t^\ell(a) = a$. Thus the A -module $\blacktriangleright (A_S)$ is free with basis 1.
- (ii) If $a \in A$, then $\alpha \cdot_S s^\ell(a) = \mu\nu(a)\alpha$. Thus the A^{op} -module $A_S \blacktriangleleft$ is free with basis 1.
- (iii) If N is a right H^ℓ -module, the left H^ℓ -module $\text{Hom}_A(\blacktriangleright (A_S), \blacktriangleright N)$ is isomorphic to ${}_S N$.
- (iv) The A^e -module $\blacktriangleright A_S \blacktriangleleft$ (defined from the right H^ℓ -module structure on A_S) is invertible.

Proof. (i) Using Proposition 3.11, we have:

$$1 \cdot_S t^\ell(a) = S(t^\ell(a))[1] \stackrel{Prop. 3.11}{=} t^r \nu(a)[1] = \epsilon [t^r \nu(a)] = a.$$

(ii) Similarly, on has: $1 \cdot_S s^\ell(a) = S(s^\ell(a))(1) = \epsilon s^r \nu(a) = \mu \nu(a)$.

(iii) The map

$$\begin{aligned} Hom_A(\blacktriangleright A_S, \blacktriangleright N) &\rightarrow {}_S N \\ \lambda &\mapsto \lambda(1) \end{aligned}$$

is an isomorphism of left H^ℓ -modules as shows the following computation. Let $\alpha \in A_S$, $h \in H^\ell$ and $\lambda \in Hom_A(\blacktriangleright A_S, \blacktriangleright N)$. Using Assertion 1 and Theorem 3.11, we have:

$$\begin{aligned} (h \cdot \lambda)(1) &= \lambda(1 \cdot_S h^{(1)})S(h^{(2)}) \\ &= \lambda[S(h^{(1)})(1)]S(h^{(2)}) \\ &= \lambda[\epsilon S(h^{(1)})]S(h^{(2)}) \\ &= \lambda[1 \cdot_S t^\ell \epsilon S(h^{(1)})]S(h^{(2)}) \\ &= \lambda(1)t^\ell \epsilon [S(h^{(1)})]S(h^{(2)}) \\ &= \lambda(1)t^\ell \epsilon [S(h)_{(2)}]S(h)_{(1)} \\ &= \lambda(1)S(h). \end{aligned}$$

(iv) Let N be a right H^ℓ -module and let $n \in N$. Denote by λ_n the element of $Hom_A(\blacktriangleright A_S, \blacktriangleright N)$ determined by $\lambda_n(1) = n$. By assertions 1 and 2, the map $(A_S) \blacktriangleleft_{A \triangleright} Hom_A(\blacktriangleright A_S, \blacktriangleright N) \rightarrow N$, $p \otimes_{A^{op}} \phi \mapsto \phi(p)$ is an isomorphism with inverse $n \mapsto 1 \otimes \lambda_n$.

We need now to check that the map $A \rightarrow Hom_A(\blacktriangleright A_S, \blacktriangleright A_S)$, $a \mapsto \{p \mapsto p \blacktriangleleft a\}$ is an isomorphism. By assertion 3, this boils down to showing that $A \rightarrow {}_S(A_S)$, $a \mapsto 1 \blacktriangleleft a$ is an isomorphism. But, this is true as $1 \blacktriangleleft a = \mu \nu(a)$. Indeed,

$$1 \blacktriangleleft a = S^2(s^\ell(a))(1) = \epsilon S^2[s^\ell(a)] = \mu \partial [S(s^\ell(a))] = \mu \nu \epsilon (s^\ell(a)) = \mu \nu(a).$$

□

We can now state *twisted Poincaré duality for full Hopf algebroids*.

Theorem 3.16. *Let (A, H^ℓ, H^r) be a Hopf algebroid over A with bijective antipode S . As in Proposition 3.11, we set $\mu = \epsilon s^r$ and $\nu = \partial s^\ell$. Assume the following:*

- (i) $\text{Ext}_{H^\ell}^i(A, H^\ell) = \{0\}$ if $i \neq d$ and set $\Lambda = \text{Ext}_{H^\ell}^d(A, H^\ell)$.
- (ii) $\blacktriangleright \text{Ext}_{H^\ell}^d(A, H^\ell)$ is a projective A -module and $\text{Ext}_{H^\ell}^d(A, H^\ell) \blacktriangleleft$ is a projective A^{op} -module.
- (iii) The left H^ℓ -module A admits a finitely generated projective resolution of finite length.

Then for all left H -modules M and all $i \in \mathbb{N}$, there is an isomorphism

$$\text{Ext}_{H^\ell}^i(A, M) \simeq \text{Tor}_{d-i}^{H^\ell}(A_S, S\Lambda \blacktriangleleft_{A \triangleright} M).$$

As an application, we find a Poincaré duality for smooth Poisson algebras. Assume that X is a C^∞ Poisson manifold, $L = \Omega^1(X)$ and M is a $V(L)$ -module. Huebschmann ([11]) has shown that for any $i \in \mathbb{N}$, the k -space $\text{Ext}_{V(\Omega^1(X))}^i(A, M)$ coincides with the i^{th} Poisson cohomology space with coefficients in M , $H_{Pois}^i(A, M)$. Also, the k -space $\text{Tor}_i^{V(\Omega^1(X))}(A_S, M)$ coincides with the i^{th} Poisson homology space with coefficients in M , $H_i^{Pois}(A, M)$.

Corollary 3.17. *Let X be a C^∞ n -dimensional Poisson manifold, $A = C^\infty(X)$ and M a left $V(\Omega^1(X))$ -module. Let S be the antipode of the (full) Hopf algebroid $V(\Omega^1(X))$ (see Examples 3.12). Then \mathcal{T} is isomorphic to ${}_S[\Lambda_A^n \Omega^1(X)^*] = {}_S[\Lambda_A^n \text{Der}(A)]$ where df acts (on the right) on $\Lambda_A^n \text{Der}(A)$ as the opposite of the Lie derivative of the Hamiltonian vector field H_f (see Remark 3.6). For all $i \in \mathbb{N}$, there is an isomorphism*

$$H_{Pois}^i(A, M) \simeq H_{n-i}^{Pois}(A, {}_S[\Lambda_A^n \text{Der}(A)] \otimes_A M).$$

Remark 3.18. This formula is proved in [8] for oriented Poisson manifolds and $M = A$ (see also [20] for polynomial algebras with quadratic Poisson structures, [30] for linear Poisson structures, [21] for general polynomial Poisson algebras).

Acknowledgment

I thank N. Kowalzig for comments on an earlier version of this text.

REFERENCES

- [1] G. Böhm, *Hopf algebroids*, Handbook of algebra, Vol. 6, Elsevier/North Holland, Amsterdam (2009), 173–236.
- [2] G. Böhm and K. Szlachányi, *Hopf algebroids with bijective antipodes: axioms, integrals, and duals*, J. Algebra **274** (2004), no. 2, 708–750.
- [3] A. Borel, P.-P. Grivel, B. Kaup, A. Haefliger, B. Malgrange, and F. Ehlers, *Algebraic D-modules*, Perspectives in Mathematics, vol. 2, Academic Press, Inc., Boston, MA, 1987.
- [4] J.-L. Brylinski, *A differential complex for Poisson manifolds*, J. of Diff. Geom., **28**, (1988), 93-114.
- [5] S. Chemla, *Poincaré duality for k - A Lie superalgebras*, Bull. Soc. Math. France **122** (1994), no. 3, 371–397.
- [6] ———, *A duality property for complex Lie algebroids*, Math. Zeitschrift **232** (1999), 367–388.
- [7] S. Chemla, F. Gavarini and N. Kowalzig *Duality features of left Hopf algebroids*, Algebras and representation theory **19**, (2016), no. 4, 913-941.
- [8] X Chen, L Liu, S Yu and J. Zheng *Batalin-Vilkovisky and gravity algebras on Poisson manifolds with semi-simple modular symmetry*, ArXiv: 2104.14099v1 [math.DG].
- [9] S. Evens, J-H Lu, A. Weinstein, *Transverse measures, the modular class and a cohomology pairing for Lie algebroid*, Quarterly Journal of Math (2) **50** (1999), 417-436.
- [10] A. Fröhlich *The Picard group of non-commutative rings, in particular of orders*, Transactions of the American Mathematical Society, volume 180 (1973), 1-45..
- [11] J. Huebschmann, *Poisson cohomology and quantization*, J. reine angew. Math. **408** (1990), 57-113.
- [12] N. Jacobsen, *Basic algebra II*, Freeman, San Francisco 1980.
- [13] L. Kadison and K. Szlachányi, *Bialgebroid actions on depth two extensions and duality*, Adv. Math. **179** (2003), no. 1, 75-121.

- [14] M. Kashiwara, *D-modules and microlocal calculus*, Translations of Mathematical Monographs, vol. 217, American Mathematical Society, Providence, RI, 2003.
- [15] Y. Kosmann-Schwarzbach, *Poisson manifolds, Lie algebroids, modular classes: a survey*, SIGMA, 4 (2008), Paper 005.
- [16] J-L. Koszul, *Crochet de Schouten-Nijenhuis et cohomologie*, E. Cartan et les mathématiciens d'aujourd'hui, Lyon 25-29 juin 1984, Astérisque hors série (1985), 257-271.
- [17] N. Kowalzig, *Hopf algebroids and their cyclic theory*, Ph. D. thesis, Universiteit Utrecht and Universiteit van Amsterdam, 2009.
- [18] N. Kowalzig and U. Krähmer, *Duality and products in algebraic (co)homology theories*, J. Algebra **323** (2010), no. 7, 2063–2081.
- [19] N. Kowalzig and H. Posthuma, *The cyclic theory of Hopf algebroids*, J. Noncomm. Geom. **5** (2011), no. 3, 423–476.
- [20] S. Launois and L. Richard, *Twisted Poincaré duality for some quadratic Poisson algebras*, Lett. Math. Phys. **79** (2007),no.2, 161-174.
- [21] J. Luo, S.-Q. Wang and Q.-S. Wu, *Twisted Poincaré duality between Poisson homology and Poisson cohomology*, J. Algebra **442** (2015), 484-505.
- [22] K.C.H Mackenzie and Ping Xu, *Lie bialgebroids and Poisson groupoids*, Duke Math. Journal **73** (1994),no. 2, 415–452.
- [23] G. Rinehart, *Differential forms on general commutative algebras*, Trans. Amer. Math. Soc. **108** (1963), 195–222.
- [24] P. Schauenburg, *Duals and doubles of quantum groupoids (\times_R -Hopf algebras)*, New trends in Hopf algebra theory (La Falda, 1999), Contemp. Math., vol. 267, Amer. Math. Soc., Providence, RI, (2000), 273–299.
- [25] M. Sweedler, *Hopf algebras*, Mathematics Lecture Note Series, W. A. Benjamin, Inc., New York, 1969.
- [26] M. Takeuchi, *Groups of algebras over $A \otimes \bar{A}$* , J. Math. Soc. Japan **29** (1977), no. 3, 459–492.
- [27] M. Van Den Bergh, *A relation between Hochschild homology and cohomology for Gorenstein rings*, Proceedings of the American Mathematical Society, **126**, No 5 (1998), 1345-1348.
- [28] P. Xu, *Quantum groupoids*, Comm. Math. Phys. **216** (2001), no. 3, 539–581.
- [29] A. Yekutieli, *Dualizing complexes over non-commutative graded algebras*, Journal of Algebra **153**, (1992), no.1, 41-84.
- [30] C. Zhu, *Twisted Poincaré duality for Poisson homology and cohomology of affine Poisson algebras*, Proc. Amer. Math. Soc., **143** (2015), no. 5, 1957-1967.

Sophie Chemla

Sorbonne Université, Université de Paris, CNRS, Institut de Mathématiques de Jussieu-Paris Rive Gauche, F-75005 Paris, France.

sophie.chemla@sorbonne-universite.fr