

# COHOMOLOGICAL CONSTRUCTION OF RELATIVE TWISTS

VALERIO TOLEDANO LAREDO

ABSTRACT. Let  $\mathfrak{g}$  be a complex, semi-simple Lie algebra,  $\mathfrak{h} \subset \mathfrak{g}$  a Cartan subalgebra and  $D$  a subdiagram of the Dynkin diagram of  $\mathfrak{g}$ . Let  $\mathfrak{g}_D \subset \mathfrak{l}_D \subseteq \mathfrak{g}$  be the corresponding semi-simple and Levi subalgebras and consider two invariant solutions  $\Phi \in (U\mathfrak{g}^{\otimes 3}[[\hbar]])^{\mathfrak{g}}$  and  $\Phi_D \in (U\mathfrak{g}_D^{\otimes 3}[[\hbar]])^{\mathfrak{g}_D}$  of the pentagon equation for  $\mathfrak{g}$  and  $\mathfrak{g}_D$  respectively. Motivated by the theory of quasi-Coxeter quasitriangular quasibialgebras [TL3], we study in this paper the existence of a *relative* twist, that is an element  $F \in (U\mathfrak{g}^{\otimes 2}[[\hbar]])^{\mathfrak{l}_D}$  such that the twist of  $\Phi$  by  $F$  is  $\Phi_D$ . Adapting the method of Donin and Shnider [DS], who treated the case of an empty  $D$ , so that  $\mathfrak{l}_D = \mathfrak{h}$  and  $\Phi_D = 1^{\otimes 3}$ , we give a cohomological construction of such an  $F$  under the assumption that  $\Phi_D$  is the image of  $\Phi$  under the generalised Harish-Chandra homomorphism  $(U\mathfrak{g}^{\otimes 3})^{\mathfrak{l}_D} \rightarrow (U\mathfrak{g}_D^{\otimes 3})^{\mathfrak{g}_D}$ . We also show that  $F$  is unique up to a gauge transformation if  $\mathfrak{l}_D$  is of corank 1 or  $F$  satisfies  $F^\Theta = F^{21}$  where  $\Theta \in \text{Aut}(\mathfrak{g})$  is an involution acting as  $-1$  on  $\mathfrak{h}$ .

## 1. INTRODUCTION

Let  $\mathfrak{g}$  be a complex, semi-simple Lie algebra,  $\mathfrak{h} \subset \mathfrak{g}$  a Cartan subalgebra and  $D_{\mathfrak{g}}$  the Dynkin diagram of  $\mathfrak{g}$  relative to a choice  $\alpha_1, \dots, \alpha_n \in \mathfrak{h}^*$  of simple roots of  $\mathfrak{g}$ . Let  $D \subseteq D_{\mathfrak{g}}$  be a subdiagram of the Dynkin diagram of  $\mathfrak{g}$  and denote by

$$\mathfrak{g}_D \subseteq \mathfrak{l}_D \subseteq \mathfrak{g}$$

the corresponding diagrammatic subalgebra, *i.e.*, the semi-simple subalgebra generated by the root vectors corresponding to the simple roots in  $D$ , and Levi subalgebra  $\mathfrak{l}_D = \mathfrak{g}_D + \mathfrak{h}$  respectively. Note that

$$\mathfrak{l}_D = \mathfrak{g}_D \oplus \mathfrak{c}_D$$

where the centre  $\mathfrak{c}_D$  of  $\mathfrak{l}_D$  is spanned by the fundamental coweights  $\lambda_j^\vee \in \mathfrak{h}$ , with  $j$  such that  $\alpha_j \notin D$ .

Let  $\hbar$  be a formal variable and consider two fixed, invariant elements

$$\Phi \in 1 + \hbar^2 (U\mathfrak{g}^{\otimes 3}[[\hbar]])^{\mathfrak{g}} \quad \text{and} \quad \Phi_D \in 1 + \hbar^2 (U\mathfrak{g}_D^{\otimes 3}[[\hbar]])^{\mathfrak{g}_D}$$

---

*Date:* June 2005.

satisfying the pentagon equation

$$\mathrm{id}^{\otimes 2} \otimes \Delta(\Psi) \cdot \Delta \otimes \mathrm{id}^{\otimes 2}(\Psi) = 1 \otimes \Psi \cdot \mathrm{id} \otimes \Delta \otimes \mathrm{id}(\Psi) \cdot \Psi \otimes 1 \quad (1.1)$$

We shall be concerned in this paper with the cohomological solution of the following *relative twist equation*

$$(\Phi)_F := 1 \otimes F \cdot \mathrm{id} \otimes \Delta(F) \cdot \Phi \cdot \Delta \otimes \mathrm{id}(F^{-1}) \cdot F^{-1} \otimes 1 = \Phi_D \quad (1.2)$$

with respect to a twist  $F$  which is invariant under the adjoint action of  $\mathfrak{l}_D$

$$F \in 1 + \hbar (U\mathfrak{g}^{\otimes 2}[[\hbar]])^{\mathfrak{l}_D}$$

Our motivation for studying (1.2) comes from the theory of quasi-Coxeter quasitriangular quasibialgebras [TL3]. These are, informally speaking, bialgebras which carry representations of both Artin's braid groups  $B_n$  and the generalised braid group  $B_{\mathfrak{g}}$  of type  $\mathfrak{g}$  on the tensor products of their finite-dimensional modules. One of the main results in [TL3] is the rigidity of quasi-Coxeter quasitriangular quasibialgebra structures on  $U\mathfrak{g}[[\hbar]]$ . In conjunction with the results of [TL4], this shows in particular that the monodromy of the Casimir connection introduced in [MTL] is described by Lusztig's quantum Weyl group operators [Lu], thus proving a conjecture formulated independently by the author [TL1, TL2] and De Concini (unpublished). The rigidity result of [TL3] depends on the one hand on Drinfeld's uniqueness theorem for quasitriangular quasibialgebra deformations of  $U\mathfrak{g}$  [Dr2] and, on the other, on the uniqueness, up to gauge transformation of solutions of (1.2) when  $\mathfrak{l}_D$  is of corank 1.

Rather than incorporating the required uniqueness result into [TL3], we decided to study the existence of solutions of (1.2) as well and present our results in a separate publication. These may in fact be of independent interest since the relevant deformation complex turns out to be a perturbation of the Chevalley–Eilenberg complex for a suitable, non-coboundary Lie algebra structure on  $\mathfrak{g}^*$ . Our method is very close to that of Donin–Shnider [DS, §3] who solved the equation (1.2) when  $D = \emptyset$ , so that  $\mathfrak{l}_D = \mathfrak{h}$  and  $\Phi_D = 1^{\otimes 3}$ , and  $\Phi$  satisfies in addition

$$\Phi^{321} = \Phi^{-1} \quad \text{and} \quad \Phi^{\Theta} = \Phi \quad (1.3)$$

where  $\Theta \in \mathrm{Aut}(\mathfrak{g})$  is an involution acting as  $-1$  on  $\mathfrak{h}$ . The possibility of laddering down, that is solving (1.2) only when  $|D_{\mathfrak{g}} \setminus D| = 1$  allows us to bypass the use of (1.3) and to construct in §5 a suitable  $F$  under the sole assumption that  $\Phi_D$  is the projection of  $\Phi$  with respect to the generalised Harish–Chandra homomorphism  $(U\mathfrak{g}^{\otimes 3})^{\mathfrak{l}_D} \rightarrow (U\mathfrak{g}_D^{\otimes 3})^{\mathfrak{g}_D}$  defined in §2. Our proof proceeds along the lines of Donin and Shnider's,

the main difference being in the cohomology theory needed to deal with secondary obstructions, which is defined and computed in §4. The uniqueness of solutions of (1.2) is obtained in §6 under the weaker assumption that the infinitesimal of  $\Phi$  projects onto that of  $\Phi_D$  and that either  $\mathfrak{l}_D$  is of corank 1 or  $F$  satisfies  $F^\Theta = F^{21}$ . Section 3 contains some standard material on the classical Yang–Baxter equations.

**Remark 1.1.** A non-cohomological proof of the existence of  $F$  may be given in the case where  $\Phi$  and  $\Phi_D$  are Lie associators by adapting Etingof and Kazhdan’s method [EK]. The latter corresponds to the case when  $\mathfrak{l}_D = \mathfrak{h}$  but can be modified by replacing the Verma modules used in [EK] by their generalised counterparts obtained by inducing from the parabolic subalgebra  $\mathfrak{p}_D \subset \mathfrak{g}$  corresponding to  $D$ .

## 2. GENERALISED HARISH–CHANDRA HOMOMORPHISMS

For each  $k \geq 1$ , we define below an algebra homomorphism

$$\pi_D^k : (U\mathfrak{g}^{\otimes k})^{\mathfrak{c}_D} \longrightarrow U\mathfrak{l}_D^{\otimes k}$$

which restricts to the identity on  $U\mathfrak{l}_D^{\otimes k}$  and is equivariant with respect to adjoint action of  $\mathfrak{l}_D$ . For  $k = 1$  and  $D = \emptyset$ ,  $\pi_D^k$  is the Harish–Chandra homomorphism  $\pi : U\mathfrak{g}^{\mathfrak{h}} \rightarrow U\mathfrak{h}$ . The definition of  $\pi_D^k$  is similar to that of  $\pi$ , see *e.g.*, [Di, §7.4.1–7.4.3] which we follow closely. Write

$$\mathfrak{g} = \mathfrak{n}_D^- \oplus \mathfrak{l}_D \oplus \mathfrak{n}_D^+$$

where the nilpotent subalgebras  $\mathfrak{n}_D^\pm$  are spanned by the roots vectors  $e_\alpha, f_\alpha$  respectively, with  $\alpha$  ranging over the roots of  $\mathfrak{g}$  not lying in the root system  $R_D$  of  $\mathfrak{g}_D$ . Set

$$I_k = (U\mathfrak{g}^{\otimes k})^{\mathfrak{c}_D} \cap \sum_{i=1}^k U\mathfrak{g}^{\otimes k} \cdot (\mathfrak{n}_D^+)_i$$

where, for  $y \in U\mathfrak{g}$ ,

$$y_i = 1^{\otimes(i-1)} \otimes y \otimes 1^{\otimes(k-i)} \in U\mathfrak{g}^{\otimes k}$$

**Proposition 2.1.**

- (i)  $I_k = (U\mathfrak{g}^{\otimes k})^{\mathfrak{c}_D} \cap \sum_{i=1}^k (\mathfrak{n}_D^-)_i \cdot U\mathfrak{g}^{\otimes k}$ .
- (ii)  $I_k$  is a two-sided ideal in  $(U\mathfrak{g}^{\otimes k})^{\mathfrak{c}_D}$  invariant under the adjoint action of  $\mathfrak{l}_D$ .
- (iii)  $(U\mathfrak{g}^{\otimes k})^{\mathfrak{c}_D} = I_k \oplus U\mathfrak{l}_D^{\otimes k}$ .

PROOF. (i) By the PBW theorem,

$$U\mathfrak{g}^{\otimes k} \cong U\mathfrak{n}_D^{-\otimes k} \otimes U\mathfrak{l}_D^{\otimes k} \otimes U\mathfrak{n}_D^{+\otimes k}$$

is spanned by the monomials

$$u(q_i^j; x; p_i^j) = f_{\beta_1,1}^{q_1^1} \cdots f_{\beta_m,1}^{q_m^1} \cdots f_{\beta_1,k}^{q_1^k} \cdots f_{\beta_m,k}^{q_m^k} \cdot x \cdot e_{\beta_1,1}^{q_1^1} \cdots e_{\beta_m,1}^{q_m^1} \cdots e_{\beta_1,k}^{q_1^k} \cdots e_{\beta_m,k}^{q_m^k}$$

where  $x \in U\mathfrak{l}_D^{\otimes k}$ ,  $\beta_1, \dots, \beta_m$  are the positive roots in  $R_{\mathfrak{g}} \setminus R_D$  and  $q_i^j, p_i^j \in \mathbb{N}$ . Let  $i^* : \mathfrak{h}^* \rightarrow \mathfrak{c}_D^*$  be the restriction map. Since  $u(q_i^j; x; p_i^j)$  has weight  $i^* \sum_{i,j} p_{i,j} \beta_j - i^* \sum_{i,j} q_{i,j} \beta_j$  for the adjoint action of  $\mathfrak{c}_D$ ,  $(U\mathfrak{g}^{\otimes k})^{\mathfrak{c}_D}$  is spanned by the  $u(q_i^j; x; p_i^j)$  such that

$$i^* \sum_{i,j} p_{i,j} \beta_j = i^* \sum_{i,j} q_{i,j} \beta_j$$

Note that, since each  $\beta_j$  restricts on  $\mathfrak{c}_D$  to a non-trivial linear combination of the fundamental coweights  $\lambda_j^\vee$ ,  $j \notin D$ , with non-negative coefficients,  $i^* \sum_{i,j} p_{i,j} \beta_j = 0$  iff  $\sum_{i,j} p_{i,j} = 0$ . It follows that

$$\begin{aligned} I_k &= \langle u(q_i^j; x; p_i^j) \in (U\mathfrak{g}^{\otimes k})^{\mathfrak{c}_D} \rangle_{\sum_{i,j} p_i^j > 0} \\ &= \langle u(q_i^j; x; p_i^j) \in (U\mathfrak{g}^{\otimes k})^{\mathfrak{c}_D} \rangle_{\sum_{i,j} q_i^j > 0} \\ &= (U\mathfrak{g}^{\otimes k})^{\mathfrak{c}_D} \bigcap \sum_{i=1}^k (\mathfrak{n}_D^-)_i \cdot U\mathfrak{g}^{\otimes k} \end{aligned}$$

as claimed. (ii)  $I_k$  is a left ideal by definition and, by (i), it is also a right ideal. It is moreover invariant under the adjoint action of  $\mathfrak{l}_D$  since  $\mathfrak{n}_D^\pm$  are. (iii) is now obvious ■

**Corollary 2.2.** *The projection  $\pi_D^k$  of  $(U\mathfrak{g}^{\otimes k})^{\mathfrak{c}_D}$  onto  $U\mathfrak{l}_D^{\otimes k}$  defined by the ideal  $I_k$  is equivariant for the adjoint action of  $\mathfrak{l}_D$  and therefore gives rise to the following commutative diagram of algebra homomorphisms*

$$\begin{array}{ccccc} (U\mathfrak{g}^{\otimes k})^{\mathfrak{c}_D} & \xrightarrow{\pi_D^k} & U\mathfrak{l}_D^{\otimes k} & \longrightarrow & U\mathfrak{g}_D^{\otimes k} \\ & & \cup & & \cup \\ & & \cup & & \cup \\ & & \cup & & \cup \\ (U\mathfrak{g}^{\otimes k})^{\mathfrak{l}_D} & \xrightarrow{\bar{\pi}_D^k} & (U\mathfrak{l}_D^{\otimes k})^{\mathfrak{l}_D} & \longrightarrow & (U\mathfrak{g}_D^{\otimes k})^{\mathfrak{g}_D} \end{array}$$

where the rightmost horizontal arrows are induced by the Lie algebra projection  $\mathfrak{l}_D \rightarrow \mathfrak{g}_D$ .

**Definition 2.3.** *We denote the composition of the horizontal arrows by  $\bar{\pi}_D^k$  and refer to  $\bar{\pi}_D^k$  or  $\pi_D^k$  as generalised Harish–Chandra homomorphisms.*

Note that  $\pi_D^k$  and  $\bar{\pi}_D^k$  are equivariant under the natural action of the symmetric group  $\mathfrak{S}_k$ . We record for later use the following two results

**Proposition 2.4.** *For any  $i, l \leq k$ ,  $x \in (U\mathfrak{g}^{\otimes k})^{\mathfrak{c}D}$  and  $y \in (U\mathfrak{g}^{\otimes l})^{\mathfrak{c}D}$ , one has*

$$\mathrm{id}^{\otimes i} \otimes \Delta \otimes \mathrm{id}^{\otimes(k-i-1)} \circ \pi_D^k(x) = \pi_D^{k+1} \circ \mathrm{id}^{\otimes i} \otimes \Delta \otimes \mathrm{id}^{\otimes(k-i-1)}(x) \quad (2.1)$$

$$1^{\otimes i} \otimes \pi_D^l(y) \otimes 1^{\otimes(k-l-i)} = \pi_D^k(1^{\otimes i} \otimes y \otimes 1^{\otimes(k-i-l)}) \quad (2.2)$$

*These identities remain valid if  $\pi_D^k, \pi_D^{k+1}$  and  $\pi_D^l$  are replaced by  $\bar{\pi}_D^k, \bar{\pi}_D^{k+1}$  and  $\bar{\pi}_D^l$  respectively.*

PROOF. Let

$$\gamma = \mathrm{id}^{\otimes i} \otimes \Delta \otimes \mathrm{id}^{\otimes(k-i-1)} : U\mathfrak{g}^{\otimes k} \rightarrow U\mathfrak{g}^{\otimes(k+1)}$$

Since  $\gamma$  is equivariant for the adjoint action of  $\mathfrak{g}$ , it maps  $(U\mathfrak{g}^{\otimes k})^{\mathfrak{c}D}$  to  $(U\mathfrak{g}^{\otimes(k+1)})^{\mathfrak{c}D}$  so that the right-hand side of (2.1) is well-defined. One readily checks that

$$\gamma(I_k) \subset I_{k+1} \quad \text{and that} \quad \gamma(U\mathfrak{U}_D^{\otimes k}) \subset U\mathfrak{U}_D^{\otimes(k+1)}$$

so that (2.1) holds. (2.2) is proved in the same way. The fact that these identities hold when  $\pi_D^j$  is replaced by  $\bar{\pi}_D^j$  throughout follows from the fact that  $\bar{\pi}_D^j = \pi^{\otimes j} \circ \pi_D^j$  where  $\pi : U\mathfrak{U}_D \rightarrow U\mathfrak{g}_D$  is a Hopf algebra homomorphism ■

**Corollary 2.5.** *Let  $d_H : U\mathfrak{g}^{\otimes k} \rightarrow U\mathfrak{g}^{\otimes(k+1)}$  be the Hochschild differential given by*

$$d_H x = 1 \otimes x + \sum_{i=1}^k (-1)^i \mathrm{id}^{\otimes(i-1)} \otimes \Delta \otimes \mathrm{id}^{\otimes(k-i)}(x) + (-1)^{k+1} x \otimes 1$$

Then,

$$d_H \circ \pi_D^k = \pi_D^{k+1} \circ d_H \quad \text{and} \quad d_H \circ \bar{\pi}_D^k = \bar{\pi}_D^{k+1} \circ d_H$$

### 3. CLASSICAL YANG–BAXTER EQUATIONS

We review below some well-known results on the classical Yang–Baxter equations due to Drinfeld [Dr1].

Define the classical Yang–Baxter map  $\mathrm{YB} : \mathfrak{g}^{\otimes 2} \otimes \mathfrak{g}^{\otimes 2} \rightarrow \mathfrak{g}^{\otimes 3}$  by

$$\mathrm{YB}(r, s) = [r^{12}, s^{13} + s^{23}] + [r^{13}, s^{23}] + [s^{12}, r^{13} + r^{23}] + [s^{13}, r^{23}]$$

Identify the exterior algebra  $\bigwedge \mathfrak{g}$  with its image in the tensor algebra  $T\mathfrak{g}$  via the antisymmetrisation map

$$X_1 \wedge \dots \wedge X_k \longrightarrow \mathrm{Alt}_k(X_1 \otimes \dots \otimes X_k) = \frac{1}{k!} \sum_{\sigma \in \mathfrak{S}_k} (-1)^{\sigma(1)} X_{\sigma(1)} \otimes \dots \otimes X_{\sigma(k)}$$

One readily checks that if  $r, s \in \bigwedge^2 \mathfrak{g}$ , then

$$\text{YB}(r, s) = 6 \text{Alt}_3[r^{12}, s^{13}] \quad (3.1)$$

**Lemma 3.1.** *If  $r = r_1 \wedge r_2, s = s_1 \wedge s_2 \in \bigwedge^2 \mathfrak{g}$ , then*

$$\text{YB}(r, s) = \frac{3}{2} \sum_{1 \leq i, j \leq 2} [r_i, s_j] \wedge r_{3-i} \wedge s_{3-j} \quad (3.2)$$

PROOF. We have

$$\begin{aligned} 4[r^{12}, s^{13}] &= [r_1, s_1] \otimes r_2 \otimes s_2 - [r_1, s_2] \otimes r_2 \otimes s_1 \\ &\quad - [r_2, s_1] \otimes r_1 \otimes s_2 + [r_2, s_2] \otimes r_1 \otimes s_1 \end{aligned}$$

Antisymmetrising both sides and using (3.1), we find (3.2) ■

Let  $(\cdot, \cdot)$  be a non-degenerate, ad-invariant, symmetric bilinear form on  $\mathfrak{g}$  and let  $\Omega = \sum_i x_i \otimes x^i$ , where  $\{x_i\}, \{x^i\}$  are dual basis of  $\mathfrak{g}$  with respect to  $(\cdot, \cdot)$ , be the corresponding symmetric, invariant tensor in  $\mathfrak{g} \otimes \mathfrak{g}$ . It is well-known that  $[\Omega_{12}, \Omega_{23}]$  lies in  $(\bigwedge^3 \mathfrak{g})^{\mathfrak{g}}$  and generates it if  $\mathfrak{g}$  is simple. Let

$$r_{\mathfrak{g}} = \sum_{\alpha > 0} \frac{(\alpha, \alpha)}{2} \cdot e_{\alpha} \wedge f_{\alpha} \in \bigwedge^2 \mathfrak{g} \quad (3.3)$$

where  $e_{\alpha} \in \mathfrak{g}_{\alpha}, f_{\alpha} \in \mathfrak{g}_{-\alpha}$  are root vectors such that  $[e_{\alpha}, f_{\alpha}] = h_{\alpha}$  so that

$$(e_{\alpha}, f_{\alpha}) = \frac{1}{2}([h_{\alpha}, e_{\alpha}], f_{\alpha}) = \frac{1}{2}(h_{\alpha}, [e_{\alpha}, f_{\alpha}]) = \frac{1}{2}(h_{\alpha}, h_{\alpha}) = \frac{2}{(\alpha, \alpha)} \quad (3.4)$$

By the following result,  $r_{\mathfrak{g}}$  is a solution of the modified classical Yang-Baxter equation (MCYBE), that is the equation

$$[r_{\mathfrak{g}}^{12}, r_{\mathfrak{g}}^{23} + r_{\mathfrak{g}}^{13}] + [r_{\mathfrak{g}}^{13}, r_{\mathfrak{g}}^{23}] \in (\bigwedge^3 \mathfrak{g})^{\mathfrak{g}} \quad (3.5)$$

**Proposition 3.2** (Drinfeld).

$$\text{YB}(r_{\mathfrak{g}}, r_{\mathfrak{g}}) = \frac{1}{2}[\Omega_{12}, \Omega_{23}]$$

**Remark 3.3.** We shall refer to  $r_{\mathfrak{g}}$  given by (3.3) as the standard (Drinfeld) solution of the MCYBE corresponding to the bilinear form  $(\cdot, \cdot)$ .

#### 4. CLASSICAL $r$ -MATRICES AND CHEVALLEY–EILENBERG COHOMOLOGY

4.1. The aim of this section is to compute the cohomology of the complex

$$((\bigwedge \mathfrak{g})^{\mathfrak{g}^D}, d) \quad \text{where} \quad d = \llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}^D}, \cdot \rrbracket$$

is given by the Schouten bracket with the difference of the standard solutions of the modified classical Yang–Baxter equations for  $\mathfrak{g}$  and  $\mathfrak{g}^D$  respectively.

The computation is carried out by identifying  $d$  with a perturbation of the Chevalley–Eilenberg differential on  $\bigwedge \mathfrak{g} = \bigwedge (\mathfrak{g}^*)^*$  induced by a suitable Lie algebra structure on  $\mathfrak{g}^*$ . When  $D = \emptyset$ , so that  $\mathfrak{l}_D = \mathfrak{h}$ , this identification is well-known and follows readily from the fact that the relevant Lie algebra structure on  $\mathfrak{g}^*$  is given in terms of the cobracket  $\delta : \mathfrak{g} \rightarrow \mathfrak{g} \wedge \mathfrak{g}$  defined by

$$\delta(X) = \text{ad}(X)r_{\mathfrak{g}} = -\llbracket r_{\mathfrak{g}}, X \rrbracket$$

When  $D \neq \emptyset$  the relevant Lie algebra structure on  $\mathfrak{g}^*$  is described in §4.4 and is not of coboundary type. We begin with a few reminders.

4.2. Recall that the Schouten bracket

$$\llbracket \cdot, \cdot \rrbracket : \bigwedge^k \mathfrak{g} \otimes \bigwedge^l \mathfrak{g} \rightarrow \bigwedge^{k+l-1} \mathfrak{g}$$

on the exterior algebra  $\bigwedge \mathfrak{g}$  is defined by

$$\begin{aligned} & \llbracket X_1 \wedge \cdots \wedge X_k, Y_1 \wedge \cdots \wedge Y_l \rrbracket \\ &= \sum_{i,j} (-1)^{i+j} [X_i, Y_j] \wedge X_1 \wedge \cdots \wedge \widehat{X}_i \wedge \cdots \wedge X_k \wedge Y_1 \wedge \cdots \wedge \widehat{Y}_j \wedge \cdots \wedge Y_l \end{aligned} \tag{4.1}$$

Since

$$\llbracket \underline{X}, \underline{Y} \rrbracket = -(-1)^{(k-1)(l-1)} \llbracket \underline{Y}, \underline{X} \rrbracket$$

for any  $\underline{X} \in \bigwedge^k \mathfrak{g}$  and  $\underline{Y} \in \bigwedge^l \mathfrak{g}$  and

$$\llbracket \underline{X}, \llbracket \underline{Y}, \underline{Z} \rrbracket \rrbracket = \llbracket \llbracket \underline{X}, \underline{Y} \rrbracket, \underline{Z} \rrbracket + (-1)^{(k-1)(l-1)} \llbracket \underline{Y}, \llbracket \underline{X}, \underline{Z} \rrbracket \rrbracket$$

for any such  $\underline{X}, \underline{Y}$  and  $\underline{Z} \in \bigwedge \mathfrak{g}$ , the Schouten bracket endows  $\bigwedge \mathfrak{g}$  with the structure of a  $\mathbb{Z}$ -graded Lie algebra provided the grading is defined by

$$\text{deg}(\bigwedge^k \mathfrak{g}) = k - 1$$

In particular, any  $r \in \bigwedge^2 \mathfrak{g}$  defines a degree 1 derivation  $d_r = \llbracket r, \cdot \rrbracket$  of  $\bigwedge \mathfrak{g}$ . Its square is readily computed from

$$d_r^2(\underline{Y}) = \llbracket r, \llbracket r, \underline{Y} \rrbracket \rrbracket = \llbracket \llbracket r, r \rrbracket, \underline{Y} \rrbracket - \llbracket r, \llbracket r, \underline{Y} \rrbracket \rrbracket = \llbracket \llbracket r, r \rrbracket, \underline{Y} \rrbracket - d_r^2(\underline{Y})$$

Since  $d_r^2$  is also an algebra derivation of  $\bigwedge \mathfrak{g}$ , and  $\llbracket \underline{X}, \underline{Y} \rrbracket = -\text{ad}(Y)\underline{X}$  for any  $\underline{X} \in \bigwedge \mathfrak{g}$  and  $Y \in \mathfrak{g}$ ,  $d_r$  is a differential if, and only if

$$\llbracket r, r \rrbracket \in \left( \bigwedge^3 \mathfrak{g} \right)^{\mathfrak{g}}$$

and therefore, by lemma 3.1, iff  $r$  is a solution of the MCYBE (3.5).

4.3. Let now

$$r_{\mathfrak{g}} \in \bigwedge^2 \mathfrak{g} \quad \text{and} \quad r_{\mathfrak{g}_D} \in \bigwedge^2 \mathfrak{g}_D$$

be solutions of the MCYBE for  $\mathfrak{g}, \mathfrak{g}_D$  respectively such that  $r_{\mathfrak{g}} - r_{\mathfrak{g}_D}$  is invariant under  $\mathfrak{g}_D$ . This is the case for example if both  $r_{\mathfrak{g}}$  and  $r_{\mathfrak{g}_D}$  are the standard solutions (3.3) of MCYBE relative to a non-degenerate, ad-invariant bilinear form  $(\cdot, \cdot)$  on  $\mathfrak{g}$  and its restriction to  $\mathfrak{g}_D$  respectively. Indeed,  $\mathfrak{n}_D^+$  and  $\mathfrak{n}_D^-$  are invariant under the adjoint action of  $\mathfrak{g}_D$  and  $(\cdot, \cdot)$  yields a  $\mathfrak{g}_D$ -equivariant identification  $(\mathfrak{n}_D^+)^* \cong \mathfrak{n}_D^-$  with respect to which

$$r_{\mathfrak{g}} - r_{\mathfrak{g}_D} = \sum_{\alpha \in R_{\mathfrak{g}}^+ \setminus R_D} \frac{(\alpha, \alpha)}{2} \cdot e_{\alpha} \wedge f_{\alpha}$$

is the image in  $\bigwedge^2(\mathfrak{n}_D^+ \oplus \mathfrak{n}_D^-)$  of

$$\text{id}_{\mathfrak{n}_D^+} \in \text{End}(\mathfrak{n}_D^+) \cong \mathfrak{n}_D^+ \otimes \mathfrak{n}_D^- \subset (\mathfrak{n}_D^+ \oplus \mathfrak{n}_D^-)^{\otimes 2}$$

under the projection  $(\mathfrak{n}_D^+ \oplus \mathfrak{n}_D^-)^{\otimes 2} \rightarrow \bigwedge^2(\mathfrak{n}_D^+ \oplus \mathfrak{n}_D^-)$ . We shall need the following simple

**Lemma 4.1.** *For any  $\underline{X} = X_1 \wedge \cdots \wedge X_k \in \bigwedge^k \mathfrak{g}$ , the following holds on  $\bigwedge \mathfrak{g}$*

$$\llbracket \underline{X}, \cdot \rrbracket = (-1)^{k-1} \cdot \sum_{i=1}^k (-1)^{i-1} e(X_1 \wedge \cdots \wedge \widehat{X}_i \wedge \cdots \wedge X_k) \cdot \text{ad}(X_i) \quad (4.2)$$

where  $e(\underline{Y})$  is exterior multiplication by  $\underline{Y}$ . In particular, if  $\underline{X} \in \bigwedge \mathfrak{g}_D$  and  $\underline{Y} \in (\bigwedge \mathfrak{g})^{\mathfrak{g}_D}$ , then  $\llbracket \underline{X}, \underline{Y} \rrbracket = 0$ .

**Proposition 4.2.** *Let  $r_{\mathfrak{g}}, r_{\mathfrak{g}_D}$  be solutions of the MCYBE for  $\mathfrak{g}, \mathfrak{g}_D$  respectively such that  $r_{\mathfrak{g}} - r_{\mathfrak{g}_D}$  is invariant under  $\mathfrak{g}_D$ . Then*

- (i)  $\llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}_D}, \cdot \rrbracket$  leaves  $(\bigwedge \mathfrak{g})^{\mathfrak{g}_D}$  invariant.



- (ii) *Its restriction to  $(\bigwedge \mathfrak{g})^{\mathfrak{g}^D}$  coincides with that of  $\llbracket r_{\mathfrak{g}}, \cdot \rrbracket$  and is therefore a differential.*
- (iii)  $\llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}^D}, r_{\mathfrak{g}} - r_{\mathfrak{g}^D} \rrbracket = \llbracket r_{\mathfrak{g}}, r_{\mathfrak{g}} \rrbracket - \llbracket r_{\mathfrak{g}^D}, r_{\mathfrak{g}^D} \rrbracket$ .

PROOF. (i) Since  $r_{\mathfrak{g}} - r_{\mathfrak{g}^D}$  is invariant under  $\mathfrak{g}^D$ , and the Schouten bracket is equivariant for the adjoint action of  $\mathfrak{g}$ ,  $\llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}^D}, \cdot \rrbracket$  leaves  $(\bigwedge \mathfrak{g})^{\mathfrak{g}^D}$  invariant. (ii) By lemma 4.1,  $\llbracket r_{\mathfrak{g}^D}, \underline{Y} \rrbracket = 0$  for any  $\underline{Y} \in (\bigwedge \mathfrak{g})^{\mathfrak{g}^D}$  so that

$$\llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}^D}, \underline{Y} \rrbracket = \llbracket r_{\mathfrak{g}}, \underline{Y} \rrbracket \quad (4.3)$$

for any such  $\underline{Y}$ . (iii) Since  $r_{\mathfrak{g}} - r_{\mathfrak{g}^D}$  is invariant under  $\mathfrak{g}^D$ , we find, by (4.3)

$$\begin{aligned} \llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}^D}, r_{\mathfrak{g}} - r_{\mathfrak{g}^D} \rrbracket &= \llbracket r_{\mathfrak{g}}, r_{\mathfrak{g}} - r_{\mathfrak{g}^D} \rrbracket \\ &= \llbracket r_{\mathfrak{g}}, r_{\mathfrak{g}} \rrbracket - \llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}^D}, r_{\mathfrak{g}^D} \rrbracket - \llbracket r_{\mathfrak{g}^D}, r_{\mathfrak{g}^D} \rrbracket = \llbracket r_{\mathfrak{g}}, r_{\mathfrak{g}} \rrbracket - \llbracket r_{\mathfrak{g}^D}, r_{\mathfrak{g}^D} \rrbracket \end{aligned}$$

as claimed  $\blacksquare$

**Remark 4.3.** Note that the proof of (iii) only uses the  $\mathfrak{g}^D$ -invariance of  $r_{\mathfrak{g}} - r_{\mathfrak{g}^D}$ . Thus, if  $r_{\mathfrak{g}} \in \bigwedge^2 \mathfrak{g}$  is a solution of the MCYBE and  $r_{\mathfrak{g}^D} \in \bigwedge^2 \mathfrak{g}^D$  is such that  $r_{\mathfrak{g}} - r_{\mathfrak{g}^D}$  is invariant under  $\mathfrak{g}^D$ , then  $r_{\mathfrak{g}^D}$  is a solution of the MCYBE for  $\mathfrak{g}^D$ . Moreover, if  $r_{\mathfrak{g}}$  is the standard solution of the MCYBE then so is  $r_{\mathfrak{g}^D}$ . Indeed, if  $\pi \in \text{End}(\bigwedge^2 \mathfrak{g})$  is the projection onto  $\mathfrak{g}^D$ -invariants, then

$$\begin{aligned} r_{\mathfrak{g}} - r_{\mathfrak{g}^D} &= \pi(r_{\mathfrak{g}} - r_{\mathfrak{g}^D}) \\ &= \pi(r_{\mathfrak{g}} - \bar{\pi}_D^2(r_{\mathfrak{g}}) + \bar{\pi}_D^2(r_{\mathfrak{g}}) - r_{\mathfrak{g}^D}) = r_{\mathfrak{g}} - \bar{\pi}_D^2(r_{\mathfrak{g}}) \end{aligned}$$

where the last equality follows from the  $\mathfrak{g}^D$ -invariance of  $r_{\mathfrak{g}} - \bar{\pi}_D^2(r_{\mathfrak{g}})$  and the fact that  $\pi(\bar{\pi}_D^2(r_{\mathfrak{g}}) - r_{\mathfrak{g}^D}) \in (\bigwedge^2 \mathfrak{g}^D)^{\mathfrak{g}^D} = \{0\}$ . Thus,  $r_{\mathfrak{g}^D} = \bar{\pi}_D^2(r_{\mathfrak{g}})$  is the standard solution of the MCYBE for  $\mathfrak{g}^D$ .

4.4. Identify  $\mathfrak{g}^*$  and  $\mathfrak{g}$  as vector spaces by using the bilinear form  $(\cdot, \cdot)$ , and endow  $\mathfrak{g}^*$  with the following Lie algebra structure

$$\mathfrak{g}^* = (\mathfrak{n}_D^+ \oplus \bar{\mathfrak{n}}_D^-) \rtimes (\mathfrak{g}^D \oplus \mathfrak{c}_D) \quad (4.4)$$

where  $\bar{\mathfrak{n}}_D^-$  is  $\mathfrak{n}_D^-$  with the opposite bracket,  $\mathfrak{g}^D$  acts on  $\mathfrak{n}_D^\pm$  by the adjoint action and  $\mathfrak{c}_D$  acts on  $\mathfrak{n}_D^\pm$  by  $\pm 1/2$  times the adjoint action. Denoting the corresponding bracket on  $\mathfrak{g}^*$  by  $[\cdot, \cdot]^*$ , we therefore have

$$\begin{aligned} [x, y]^* &= [x_D, y_D] + [x_D, y_+ + y_-] + [x_+ + x_-, y_D] \\ &\quad + [x_+, y_+] - [x_-, y_-] + \frac{1}{2}[x_0, y_+ - y_-] + \frac{1}{2}[x_+ - x_-, y_0] \end{aligned} \quad (4.5)$$

where,  $z_D \in \mathfrak{g}^D$ ,  $z_\pm \in \mathfrak{n}_D^\pm$  and  $z_0 \in \mathfrak{c}_D$  are the components of  $z \in \mathfrak{g}^*$  corresponding to the decomposition (4.4). Thus,  $\mathfrak{g}^D$  is a Lie subalgebra

of  $\mathfrak{g}^*$  and its coadjoint action on  $(\mathfrak{g}^*)^* = \mathfrak{g}$  coincides with its adjoint action on  $\mathfrak{g}$ .

Let now  $\delta \in \text{End}(\bigwedge \mathfrak{g})$  be the differential obtained by regarding  $\bigwedge \mathfrak{g}$  as the Chevalley–Eilenberg complex of  $\mathfrak{g}^*$ . The following result identifies  $\llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}_D}, \cdot \rrbracket$  with a perturbation of  $\delta$ .

**Theorem 4.4.** *If  $r_{\mathfrak{g}}, r_{\mathfrak{g}_D}$  are the standard solutions of the MCYBE for  $\mathfrak{g}, \mathfrak{g}_D$  respectively, then the following holds on  $\bigwedge \mathfrak{g}$*

$$\llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}_D}, \cdot \rrbracket = 2\delta + e(v_i) \cdot [\text{ad}(v_i) \cdot (1 + 2P_+)]^\wedge \quad (4.6)$$

where  $\{v_i\}, \{v^i\}$  are basis of  $\mathfrak{g}_D$  dual with respect to  $(\cdot, \cdot)$ ,  $P_+ : \mathfrak{g} \rightarrow \mathfrak{n}_D^+$  is the projection corresponding to the decomposition (4.4) and  $T \in \mathfrak{gl}(\mathfrak{g}) \rightarrow T^\wedge \in \mathfrak{gl}(\bigwedge \mathfrak{g})$  is the Lie algebra homomorphism given by

$$T^\wedge X_1 \wedge \dots \wedge X_k = \sum_{i=1}^k X_1 \wedge \dots \wedge TX_i \wedge \dots \wedge X_k$$

PROOF. It is sufficient to check (4.6) on elements of  $\mathfrak{g} \subset \bigwedge \mathfrak{g}$  since both sides are degree 1 algebra derivations of  $\bigwedge \mathfrak{g}$ . In turn, it is easier to check that the transposes of both sides coincide as maps  $\bigwedge^2 \mathfrak{g} \rightarrow \mathfrak{g}$ . By definition,  $\delta^t = [\cdot, \cdot]^*$ . Since  $e(v)^t = \iota(v)$  where  $\iota(v)$  is the contraction operator defined by

$$\iota(v) Y_1 \wedge \dots \wedge Y_l = \sum_{i=1}^l (-1)^{i-1} (v, Y_i) Y_1 \wedge \dots \wedge \widehat{Y}_i \wedge \dots \wedge Y_l$$

and  $\text{ad}(X)^t = -\text{ad}(X)$  for any  $X \in \mathfrak{g}$ , we find, using (4.2)

$$\llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}_D}, \cdot \rrbracket^t = \sum_{\alpha \in R_{\mathfrak{g}}^+ \setminus R_D} \frac{(\alpha, \alpha)}{2} (\text{ad}(e_\alpha) \iota(f_\alpha) - \text{ad}(f_\alpha) \iota(e_\alpha))$$

which, applied to  $u \wedge v \in \bigwedge^2 \mathfrak{g}$  yields

$$\begin{aligned} & \sum_{\alpha \in R_{\mathfrak{g}}^+ \setminus R_D} \frac{(\alpha, \alpha)}{2} ((f_\alpha, u)[e_\alpha, v] - (f_\alpha, v)[e_\alpha, u] - (e_\alpha, u)[f_\alpha, v] + (e_\alpha, v)[f_\alpha, u]) \\ &= [u_+, v] + [u, v_+] - [u_-, v] - [u, v_-] \\ &= 2[u_+, v_+] - 2[u_-, v_-] + [u_0, v_+ - v_-] + [u_+ - u_-, v_0] \\ & \quad + [u_+, v_D] + [u_D, v_+] - [u_-, v_D] - [u_D, v_-] \end{aligned}$$

Comparing with (4.5), we see that this is equal to

$$\begin{aligned} & 2[u, v]^* - 2[u_D, v_D] - [u_D, v_+] - [u_+, v_D] - 3[u_D, v_-] - 3[u_-, v_D] \\ &= 2[u, v]^* - [u_D, v] - [u, v_D] - 2[u_D, v_-] - 2[u_-, v_D] \\ &= 2[u, v]^* - (1 + 2P_-)([u_D, v] + [u, v_D]) \end{aligned}$$

where  $P_-$  is the projection onto  $\mathfrak{n}_D^-$  which commutes with the adjoint action of  $\mathfrak{g}_D$ . Noting that, for  $x, y \in \mathfrak{g}$ , one has

$$\text{ad}(v^i)\iota(v_i)x \wedge y = (v_i, x)[v^i, y] - (v_i, y)[v^i, x] = [x_D, y] + [x, y_D]$$

we therefore find

$$\llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}_D}, \cdot \rrbracket^t = 2[\cdot, \cdot]^* - (1 + 2P_-)\text{ad}(v_i)\iota(v^i)$$

which yields (4.6) since  $P_-^t = P_+$  ■

4.5. Note that  $(\bigwedge \mathfrak{g})^{\iota_D}$  is a subcomplex of  $((\bigwedge \mathfrak{g})^{\mathfrak{g}_D}, \llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}_D}, \cdot \rrbracket)$  since  $r_{\mathfrak{g}} - r_{\mathfrak{g}_D}$  is of weight zero. Note also that the restriction of  $\llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}_D}, \cdot \rrbracket$  to

$$(\bigwedge \mathfrak{l}_D)^{\iota_D} = (\bigwedge \mathfrak{g}_D)^{\mathfrak{g}_D} \widehat{\otimes} \bigwedge \mathfrak{c}_D \subset (\bigwedge \mathfrak{g})^{\iota_D}$$

is zero since  $r_{\mathfrak{g}} - r_{\mathfrak{g}_D}$  is invariant under  $\mathfrak{l}_D$ .

**Theorem 4.5.** *If  $r_{\mathfrak{g}}, r_{\mathfrak{g}_D}$  are the standard solutions of the MCYBE for  $\mathfrak{g}, \mathfrak{g}_D$  respectively, the inclusions*

$$\left( (\bigwedge \mathfrak{l}_D)^{\iota_D}, 0 \right) \longrightarrow \left( (\bigwedge \mathfrak{g})^{\iota_D}, \llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}_D}, \cdot \rrbracket \right) \longrightarrow \left( (\bigwedge \mathfrak{g})^{\mathfrak{g}_D}, \llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}_D}, \cdot \rrbracket \right)$$

are quasi-isomorphisms.

PROOF. Denote  $\llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}_D}, \cdot \rrbracket$  by  $d$ . It is sufficient to find an  $\mathfrak{l}_D$ -equivariant, diagonalisable operator  $C \in \text{End}(\bigwedge \mathfrak{g})$  with kernel  $\bigwedge \mathfrak{l}_D$  and an  $\mathfrak{l}_D$ -equivariant homotopy  $h \in \text{End}(\bigwedge \mathfrak{g})$  such  $dh + hd = C$ . Noting that  $\mathfrak{c}_D \subset \mathfrak{g}^*$  acts on  $\bigwedge \mathfrak{g}$  via the coadjoint action with non-negative weights only so that the corresponding subspace of invariants is precisely  $\bigwedge \mathfrak{l}_D$ , we see that a suitable  $C$  is given by the Casimir operator

$$C = \text{ad}^*(t_i) \text{ad}^*(t^i)$$

where  $\text{ad}^*$  is the coadjoint action of  $\mathfrak{g}^*$  on  $\bigwedge \mathfrak{g}$  and  $\{t_i\}, \{t^i\}$  are dual basis of  $\mathfrak{c}_D$  with respect to  $(\cdot, \cdot)$ . We claim that

$$h = \text{ad}^*(t_i)\iota(t^i)$$

satisfies  $dh + hd = 2C$ . It is well-known that  $h$  satisfies  $\delta h + h\delta = C$ , where  $\delta \in \text{End}(\bigwedge \mathfrak{g})$  is the Chevalley–Eilenberg differential. Indeed,

$$\begin{aligned} \delta h + h\delta &= \delta \text{ad}^*(t_i)\iota(t^i) + \text{ad}^*(t_i)\iota(t^i)\delta \\ &= \text{ad}^*(t_i)(\delta \iota(t^i) + \iota(t^i)\delta) \\ &= \text{ad}^*(t_i) \text{ad}^*(t^i) \end{aligned}$$

where we have used the fact that  $\delta$  is equivariant for  $\text{ad}^*$  and the identity  $\delta\iota(X) + \iota(X)\delta = \text{ad}^*(X)$ ,  $X \in \mathfrak{g}^*$ . By theorem 4.4, it therefore suffices to show that  $h$  anticommutes with

$$k = e(v_j) \cdot [(1 + 2P_+) \cdot \text{ad}(v^j)]^\wedge$$

Bearing in mind the following identities for  $X, Y \in \mathfrak{g}$  and  $T \in \mathfrak{gl}(\mathfrak{g})$

$$\begin{aligned} \iota(X)e(Y) + e(Y)\iota(X) &= (X, Y), \\ [T^\wedge, \iota(X)] &= \iota(TX) \quad \text{and} \quad [T^\wedge, e(Y)] = e(TY) \end{aligned}$$

and the fact that  $\text{ad}^*(X)^\wedge = \text{ad}^*(X)$  for any  $X \in \mathfrak{g}^*$ , we find

$$\begin{aligned} kh &= e(v_j) \cdot [(1 + 2P_+) \cdot \text{ad}(v^j)]^\wedge \cdot \text{ad}^*(t_i)\iota(t^i) \\ &= \text{ad}^*(t_i)e(v_j) \cdot [(1 + 2P_+) \cdot \text{ad}(v^j)]^\wedge \cdot \iota(t^i) \\ &= -\text{ad}^*(t_i)\iota(t^i)e(v_j) \cdot [(1 + 2P_+) \cdot \text{ad}(v^j)]^\wedge \end{aligned}$$

as claimed ■

Since  $(\bigwedge^i \mathfrak{g}_D)^{\mathfrak{g}_D} = 0$  for  $i = 1, 2$ , we obtain in particular the following

**Corollary 4.6.**

$$\begin{aligned} H^1((\bigwedge \mathfrak{g})^{\mathfrak{g}_D}; \llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}_D}, \cdot \rrbracket) &\cong \mathfrak{c}_D \\ H^2((\bigwedge \mathfrak{g})^{\mathfrak{g}_D}; \llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}_D}, \cdot \rrbracket) &\cong \bigwedge^2 \mathfrak{c}_D \\ H^3((\bigwedge \mathfrak{g})^{\mathfrak{g}_D}; \llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}_D}, \cdot \rrbracket) &\cong \bigwedge^3 \mathfrak{c}_D \oplus (\bigwedge^3 \mathfrak{g}_D)^{\mathfrak{g}_D} \end{aligned}$$

## 5. EXISTENCE OF TWISTS

Let

$$\Phi \in 1 + \hbar^2 (U\mathfrak{g}^{\otimes 3} \llbracket \hbar \rrbracket)^{\mathfrak{g}}$$

be a solution of the pentagon equation (1.1). We shall need to assume that  $\Phi$  is non-degenerate in the sense defined below. Write

$$\Phi = 1 + \hbar^2 \varphi \quad \text{mod } \hbar^3 \quad \text{where } \varphi \in (U\mathfrak{g}^{\otimes 3})^{\mathfrak{g}}$$

Taking the coefficient of  $\hbar^2$  in the pentagon relation for  $\Phi$ , we find that  $d_H \varphi = 0$  where  $d_H$  is the Hochschild differential. Thus

$$\text{Alt}_3(\varphi) \in (\bigwedge^3 \mathfrak{g})^{\mathfrak{g}} = \bigoplus_i^3 (\bigwedge \mathfrak{g}_i)^{\mathfrak{g}_i} \quad (5.1)$$

where  $\mathfrak{g}_i$  are the simple factors of  $\mathfrak{g}$ .

**Definition 5.1.**  $\Phi$  is a non-degenerate solution of the pentagon equation if the components of  $\text{Alt}_3(\varphi)$  along the decomposition (5.1) are all non-zero.

Since each  $(\bigwedge^3 \mathfrak{g}_i)^{\mathfrak{g}_i}$  is one-dimensional and generated by  $[\Omega_{12}^i, \Omega_{23}^i]$ , where  $\Omega^i \in (\mathfrak{g}_i \otimes \mathfrak{g}_i)^{\mathfrak{g}_i}$  is the symmetric element corresponding to the Killing form of  $\mathfrak{g}_i$ ,  $\Phi$  is a non-degenerate solution of the pentagon equation iff

$$\text{Alt}_3(\varphi) = \frac{1}{6}[\Omega_{12}, \Omega_{23}] \quad (5.2)$$

where  $\Omega \in (\mathfrak{g} \otimes \mathfrak{g})^{\mathfrak{g}}$  corresponds to a non-degenerate, ad-invariant, symmetric bilinear form  $(\cdot, \cdot)$  on  $\mathfrak{g}$ .

Let now  $D \subset D_{\mathfrak{g}}$  be a subdiagram, and set

$$\Phi_D = \bar{\pi}_D^3(\Phi) \in 1 + \hbar^2 (U\mathfrak{g}_D^{\otimes 3}[[\hbar]])^{\mathfrak{g}_D}$$

where  $\bar{\pi}_D^3$  is the generalised Harish–Chandra homomorphism defined in §2. By proposition 2.4,  $\Phi_D$  satisfies the pentagon equation and may therefore be regarded as an associator for  $\mathfrak{g}_D$ . Note that  $\Phi_D$  is non-degenerate if  $\Phi$  is.

**Theorem 5.2.** *If  $\Phi$  is non-degenerate, there exists a twist*

$$F \in 1 + \hbar (U\mathfrak{g}^{\otimes 2}[[\hbar]])^{\mathfrak{g}_D}$$

such that

$$(\Phi)_F = \Phi_D \quad \text{and} \quad \bar{\pi}_D^2(F) = 1 \otimes 1 \quad (5.3)$$

Modulo  $\hbar^2$ , one has

$$F = 1^{\otimes 2} + \hbar(r_{\mathfrak{g}} - r_{\mathfrak{g}_D})$$

where  $r_{\mathfrak{g}}, r_{\mathfrak{g}_D}$  are the standard solutions of the MCYBE for  $\mathfrak{g}$  and  $\mathfrak{g}_D$  corresponding to  $(\cdot, \cdot)$ . If  $\Phi$  satisfies in addition

$$\Phi^{321} = \Phi^{-1} \quad \text{and} \quad \Phi^{\Theta} = \Phi \quad (5.4)$$

where  $\Theta \in \text{Aut}(\mathfrak{g})$  is an involution acting as  $-1$  on  $\mathfrak{h}$ , then  $F$  may be chosen such that

$$F^{\Theta} = F^{21} \quad (5.5)$$

The proof of theorem 5.2 is given in 5.1–5.6. It closely follows the argument of Donin–Shnider [DS, §3] where theorem 5.2 is proved, under the additional assumption (5.4), in the case  $D = \emptyset$ . The reader familiar with Donin and Shnider’s argument will readily recognize that the only relevant difference is that the cohomology group

$$H^3(\bigwedge \mathfrak{g}; [[r_{\mathfrak{g}}, \cdot]]) \cong \bigwedge^3 \mathfrak{h} \quad (5.6)$$

which governs the secondary obstructions theory in [DS] is replaced by the group

$$H^3((\bigwedge \mathfrak{g})^{\mathfrak{g}D}; \llbracket r_{\mathfrak{g}} - r_{\mathfrak{g}D}, \cdot \rrbracket) \cong \bigwedge^3 \mathfrak{c}_D \oplus \bigoplus^3 (\bigwedge^3 \mathfrak{g}_D)^{\mathfrak{g}D} \quad (5.7)$$

which was computed in §4. Another significant difference is that the possibility of laddering down from  $D_{\mathfrak{g}}$  to  $D$  through intermediate diagrams as explained in §5.1, allows in effect to assume that  $\mathfrak{c}_D$  is at most two-dimensional, thus killing the first component of the secondary obstruction in (5.7) and rendering the assumption (5.4) unnecessary to prove the existence of  $F$ .

5.1. Note first that we may assume that  $|D_{\mathfrak{g}} \setminus D| \leq 2$ . Indeed, assume theorem 5.2 proved in this case and let

$$D_{\mathfrak{g}} = D_1 \supset D_2 \supset \cdots \supset D_{m-1} \supset D_m = D$$

be a nested chain of diagrams such that  $|D_i \setminus D_{i+1}| \leq 2$ . For any pair  $D'' \subseteq D' \subseteq D_{\mathfrak{g}}$ , denote by  $\mathfrak{c}_{D'', D'} \subset \mathfrak{h}$  the span of the fundamental coweights  $\lambda_j^{\vee}$ ,  $j \in D' \setminus D''$  and by

$$\bar{\pi}_{D'', D'}^k : (U \mathfrak{g}_{D'}^{\otimes k})^{\mathfrak{c}_{D'', D'}} \rightarrow U \mathfrak{g}_{D''}^{\otimes k}$$

the corresponding generalised Harish–Chandra homomorphism. Set  $\Phi_1 = \Phi$  and, for  $i = 1 \dots m-1$

$$\Phi_{i+1} = \bar{\pi}_{D_{i+1}, D_i}^3(\Phi_i) = \bar{\pi}_{D_{i+1}, D_{\mathfrak{g}}}^3(\Phi)$$

so that  $\Phi_m = \Phi_D$ . Let

$$F_i \in 1 + \hbar(U \mathfrak{g}_{D_i}^{\otimes 2}[[\hbar]])^{\mathfrak{g}D_{i+1}}$$

be such that

$$(\Phi_i)_{F_i} = \Phi_{i+1} \quad \text{and} \quad \bar{\pi}_{D_{i+1}, D_i}^2(F_i) = 1^{\otimes 2}$$

then

$$F = F_{m-1} \cdots F_1$$

is readily seen to satisfy (5.3). Note that if  $\Phi$  satisfies in addition (5.4) then so does each  $\Phi_i$  since  $\bar{\pi}_D^k$  is equivariant for the action of the symmetric group  $\mathfrak{S}_k$  and of  $\Theta$ . In that case, choosing each  $F_i$  such that  $F_i^{\Theta} = F_i^{21}$  yields a twist  $F$  which satisfies  $F^{\Theta} = F^{21}$ .

**Remark 5.3.** The assumption  $|D_{\mathfrak{g}} \setminus D| \leq 2$  will only be used from §5.5 onwards.

5.2. We begin by solving the equation (5.3) mod  $\hbar^2$ . Let  $f \in (U\mathfrak{g}^{\otimes 2})^{\wr_D}$  and set  $F = 1 + \hbar f$ . Since  $\Phi$  and  $\Phi_D$  are equal to 1 mod  $\hbar^2$ , the coefficient of  $\hbar$  in  $(\Phi)_F - \Phi_D$  is

$$1 \otimes f + \text{id} \otimes \Delta(f) - \Delta \otimes \text{id}(f) - f \otimes 1 = d_H f$$

Thus,  $F$  is a solution of (5.3) mod  $\hbar^2$  if, and only if,  $f$  is a Hochschild 2-cocycle such that  $\pi_D^2 f = 0$ .

5.3. Let now  $n \geq 1$  and let

$$F = 1 + \hbar f + \cdots + \hbar^n f_n \in 1 + \hbar(U\mathfrak{g}^{\otimes 2}[[\hbar]])^{\wr_D}$$

be a solution of (5.3) mod  $\hbar^{n+1}$ . We shall derive below a necessary and sufficient condition for (5.3) to possess a solution mod  $\hbar^{n+2}$  of the form  $\tilde{F} = F + \hbar^{n+1} f_{n+1}$  where

$$f_{n+1} \in (U\mathfrak{g}^{\otimes 2})^{\wr_D} \quad \text{satisfies} \quad \pi_D^2(f_{n+1}) = 0$$

Define  $\xi \in U\mathfrak{g}^{\otimes 3}$  by

$$1 \otimes F \cdot \text{id} \otimes \Delta(F) \cdot \Phi - \Phi_D \cdot F \otimes 1 \cdot \Delta \otimes \text{id}(F) = \hbar^{n+1} \xi \quad \text{mod } \hbar^{n+2} \quad (5.8)$$

Then,  $\tilde{F}$  is a solution of (5.3) mod  $\hbar^{n+2}$  if, and only if

$$d_H f_{n+1} = -\xi \quad (5.9)$$

**Lemma 5.4.** *The element  $\xi$  is invariant under  $\wr_D$  and satisfies*

$$d_H \xi = 0 \quad \text{and} \quad \pi_D^3(\xi) = 0$$

PROOF. The invariance of  $\xi$  under  $\wr_D$  follows from that of  $\Phi, \Phi_D$  and  $F$ . Since  $F = 1$  mod  $\hbar$ ,

$$\hbar^{n+1} \xi = (\Phi)_F - \Phi_D \quad \text{mod } \hbar^{n+2}$$

Since  $F$  is invariant under  $\mathfrak{g}_D$ , the restriction of

$$\Delta_F(\cdot) = F \cdot \Delta(\cdot) \cdot F^{-1}$$

to  $U\mathfrak{g}_D$  is equal to  $\Delta$  and  $\Phi_D$  satisfies the pentagon equation with respect to  $\Delta_F$ . Since this is also the case of  $(\Phi)_F$ , we find, working mod  $\hbar^{n+2}$ , that

$$0 = \text{Pent}_{\Delta_F}((\Phi)_F) = \text{Pent}_{\Delta_F}(\Phi_D) + \hbar^{n+1} d_H \xi = \hbar^{n+1} d_H \xi$$

where, for any  $\Psi \in U\mathfrak{g}^{\otimes 3}$  and map  $\tilde{\Delta} : U\mathfrak{g} \rightarrow U\mathfrak{g}^{\otimes 2}$ ,

$$\text{Pent}_{\tilde{\Delta}}(\Psi) = 1 \otimes \Psi \cdot \text{id} \otimes \tilde{\Delta} \otimes \text{id}(\Psi) \cdot \Psi \otimes 1 - \text{id}^{\otimes 2} \otimes \tilde{\Delta}(\Psi) \cdot \tilde{\Delta} \otimes \text{id}^{\otimes 2}(\Psi)$$

Finally, from  $\pi_D^2(F) = 1^{\otimes 2}$  and  $\Phi_D = \pi_D^3(\Phi)$ , we get, using proposition 2.4 that

$$\hbar^{n+1} \pi_D^3 \xi = (\pi_D^3(\Phi))_{\pi_D^2(F)} - \Phi_D = 0$$

■

**Lemma 5.5.** *If  $\Phi$  and  $F$  satisfy (5.4) and (5.5) respectively, then*

$$\xi^\Theta = -\xi^{321} \quad (5.10)$$

PROOF. We have, working mod  $\hbar^{n+2}$ ,

$$\begin{aligned} \Phi_D^\Theta + \hbar^{n+1}\xi^\Theta &= 1 \otimes F^{21} \cdot \text{id} \otimes \Delta(F^{21}) \cdot \Phi \cdot \Delta \otimes \text{id}((F^{21})^{-1}) \otimes (F^{21})^{-1} \otimes 1 \\ &= (F \otimes 1 \cdot \Delta \otimes \text{id}(F) \cdot \Phi^{321} \cdot \text{id} \otimes \Delta(F^{-1}) \cdot 1 \otimes F^{-1})^{321} \\ &= ((1 \otimes F \cdot \text{id} \otimes \Delta(F) \cdot \Phi \cdot \Delta \otimes \text{id}(F^{-1}) \cdot F^{-1} \otimes 1)^{-1})^{321} \\ &= ((\Phi_D + \hbar^{n+1}\xi)^{-1})^{321} \\ &= (\Phi_D^{-1} - \hbar^{n+1}\xi)^{321} \end{aligned}$$

whence (5.10) since  $\Phi_D = \bar{\pi}_D^3(\Phi)$  satisfies (5.4) ■

**Corollary 5.6.** *The twist  $F$  may be extended to a solution of (5.3) mod  $\hbar^{n+2}$  if, and only if  $\text{Alt}_3 \xi = 0$ . If in addition  $\Phi, F$  satisfy (5.4) and (5.5) respectively, the extension may be chosen so as to satisfy (5.5) too.*

PROOF. If  $\text{Alt}_3 \xi = 0$ , then  $\xi = d_H g$  for some  $g \in U\mathfrak{g}^{\otimes 2}$  which may be chosen invariant under  $\mathfrak{l}_D$ . By corollary 2.5, we have

$$0 = \bar{\pi}_D^3 \xi = \bar{\pi}_D^3 d_H g = d_H \bar{\pi}_D^2 g$$

so that, setting  $f_{n+1} = -(g - \bar{\pi}_D^2(g))$  we have

$$d_H f_{n+1} = -\xi \quad \text{and} \quad \bar{\pi}_D^2 f_{n+1} = 0$$

and  $F + \hbar^{n+1} f_{n+1}$  is a solution of (5.3) mod  $\hbar^{n+2}$ . If  $\Phi, F$  satisfy (5.4) and (5.5) respectively, then, by lemma 5.5

$$d_H f_{n+1}^\Theta = -\xi^\Theta = \xi^{321} = -(d_H f_{n+1})^{321} = d_H f_{n+1}^{21}$$

so that  $f'_{n+1} = 1/2(f_{n+1} + (f_{n+1}^{21})^\Theta)$  satisfies

$$d_H f'_{n+1} = \xi, \quad \bar{\pi}_D^2 f'_{n+1} = 0 \quad \text{and} \quad (f'_{n+1})^\Theta = (f'_{n+1})^{21}$$

and  $F + \hbar^{n+1} f'_{n+1}$  solves (5.3) mod  $\hbar^{n+2}$  and satisfies (5.5) ■

5.4. We consider first the case  $n = 1$  so that  $F = 1 + \hbar f$  where  $f \in (U\mathfrak{g}^{\otimes 2})^{\mathfrak{l}_D}$  is a Hochschild 2-cocycle such that  $\bar{\pi}_D^2(f) = 0$ . By lemma 5.9, adding a 2-coboundary to  $f$  does not affect the extendability of  $F$  to a solution mod  $\hbar^3$ . We may therefore assume that  $f \in (\wedge^2 \mathfrak{g})^{\mathfrak{l}_D}$ . In this case, since  $\Phi, \Phi_D$  are equal to  $1^{\otimes 3}$  mod  $\hbar^2$ , we get

$$\xi = \varphi - \varphi_D + f^{23}(f^{12} + f^{13}) - f^{12}(f^{13} + f^{23})$$

where  $\Phi = 1 + \hbar^2 \varphi$  mod  $\hbar^3$  and  $\varphi_D = \bar{\pi}_D^3 \varphi$ . Thus,  $F$  extends to a solution mod  $\hbar^3$  if, and only if,

$$\text{Alt}_3(\varphi) - \text{Alt}_3(\varphi_D) = \text{Alt}_3(f^{12}(f^{13} + f^{23}) - f^{23}(f^{12} + f^{13}))$$



We shall need the following

**Lemma 5.7.** *For any  $f, \chi \in \bigwedge^2 \mathfrak{g}$ , one has*

$$\begin{aligned} & \text{Alt}_3(f^{12}(\chi^{13} + \chi^{23}) + \chi^{12}(f^{13} + f^{23}) - f^{23}(\chi^{12} + \chi^{13}) - \chi^{23}(f^{12} + f^{13})) \\ &= \llbracket f, \chi \rrbracket \end{aligned} \tag{5.11}$$

where  $\llbracket \cdot, \cdot \rrbracket$  is the Schouten bracket (4.1).

PROOF. Since

$$\begin{aligned} (f^{12}(\chi^{13} + \chi^{23}))^{(13)} &= f^{23}(\chi^{13} + \chi^{12}), \\ (f^{12}\chi^{23})^{(12)} &= -f^{12}\chi^{13} \quad \text{and} \quad (\chi^{12}f^{13})^{(23)} = \chi^{13}f^{12} \end{aligned}$$

the left-hand side of (5.11) is equal to

$$\begin{aligned} 2 \text{Alt}_3(f^{12}(\chi^{13} + \chi^{23}) + \chi^{12}(f^{13} + f^{23})) &= 4 \text{Alt}_3(f^{12}\chi^{13} + \chi^{12}f^{13}) \\ &= 4 \text{Alt}_3(\llbracket f^{12}, \chi^{13} \rrbracket) \\ &= \frac{2}{3} \text{YB}(f, \chi) \\ &= \llbracket f, \chi \rrbracket \end{aligned}$$

where we used (3.1) and lemma 3.1 ■

**Corollary 5.8.** *Let  $f \in (\bigwedge^2 \mathfrak{g})^{\text{td}}$  be such that  $\bar{\pi}_D^2(f) = 0$ . Then, the twist*

$$F = 1 + \hbar f$$

*extends to a solution  $\tilde{F}$  of (5.3) mod  $\hbar^3$  if, and only if,*

$$\frac{1}{2} \llbracket f, f \rrbracket = \text{Alt}_3 \varphi - \text{Alt}_3 \varphi_D$$

*In that case, and provided (5.4) holds,  $\tilde{F}$  may be chosen so as to satisfy (5.5).*

Let  $(\cdot, \cdot)$  be the non-degenerate, ad-invariant, symmetric bilinear form on  $\mathfrak{g}$  such that

$$\text{Alt}_3(\varphi) = \frac{1}{6} [\Omega_{12}, \Omega_{23}] \tag{5.12}$$

and  $r_{\mathfrak{g}}, r_{\mathfrak{g}_D}$  be the standard solutions of the MCYBE determined by  $(\cdot, \cdot)$  and its restriction to  $\mathfrak{g}_D$  respectively so that  $\bar{\pi}_D^2(r_{\mathfrak{g}}) = r_{\mathfrak{g}_D}$ . We henceforth set

$$f = r_{\mathfrak{g}} - r_{\mathfrak{g}_D} \in (\bigwedge^2 \mathfrak{g})^{\text{td}} \tag{5.13}$$

By corollary 5.8,  $F = 1 + \hbar f$  extends to a solution of (5.3) mod  $\hbar^3$  which, in addition, satisfies (5.5) if (5.4) holds. Indeed, by proposition 4.2, lemma 3.1 and proposition 3.2, we have

$$\begin{aligned} \llbracket f, f \rrbracket &= \llbracket r_{\mathfrak{g}}, r_{\mathfrak{g}} \rrbracket - \llbracket r_{\mathfrak{g}_D}, r_{\mathfrak{g}_D} \rrbracket \\ &= \frac{2}{3} (\text{YB}(r_{\mathfrak{g}}, r_{\mathfrak{g}}) - \text{YB}(r_{\mathfrak{g}_D}, r_{\mathfrak{g}_D})) = \frac{1}{3} ([\Omega_{12}, \Omega_{23}] - [\Omega_{12}^D, \Omega_{23}^D]) \end{aligned}$$

5.5. Assume now  $n \geq 2$  and let

$$F = 1 + \hbar f + \hbar^2 f_2 + \cdots + \hbar^n f_n$$

be a solution of (5.3) mod  $\hbar^{n+1}$ . Let  $\xi = \xi(f; f_2, \dots, f_n) \in U\mathfrak{g}^{\otimes 3}$  be given by (5.8). By §5.3,  $\xi$  is a Hochschild 3-cocycle and  $F$  extends to a solution mod  $\hbar^{n+2}$  if, and only if  $\xi$  is a coboundary. This, however need not be the case. We note none-the-less that if  $\chi \in (U\mathfrak{g}^{\otimes 2})^{\mathfrak{l}_D}$  satisfies

$$d_H \chi = 0 \quad \text{and} \quad \bar{\pi}_D^2(\chi) = 0$$

then  $F + \hbar^n \chi$  is also a solution of (5.3) mod  $\hbar^{n+1}$  which could admit an extension mod  $\hbar^{n+2}$ . By the following result, the extendability of  $F + \hbar^n \chi$  only depends upon the Hochschild cohomology class of  $\chi$ .

**Lemma 5.9.** *If  $\chi$  is a Hochschild 2-coboundary, then the twist  $F + \hbar^n \chi$  can be extended to a solution mod  $\hbar^{n+2}$  of (5.3) if, and only if  $F$  can.*

PROOF. It suffices to prove one implication since  $F = (F + \hbar^n \chi) - \hbar^n \chi$ . Write  $\chi = d_H g$  with  $g \in U\mathfrak{g}^{\mathfrak{l}_D}$ . By corollary 2.5,

$$0 = \bar{\pi}_D^2(\chi) = d_H \bar{\pi}_D^1 g$$

so that  $\chi = d_H g'$  where  $g' = (1 - \bar{\pi}_D^1)g$  is invariant under  $\mathfrak{l}_D$  and lies in the kernel of  $\bar{\pi}_D^1$ . Let  $\tilde{F} = F + \hbar^{n+1} f_{n+1}$  be a solution mod  $\hbar^{n+2}$  of (5.3). Then

$$\tilde{F}' = u^{\otimes 2} \cdot \tilde{F} \cdot \Delta(u)^{-1}$$

where  $u = 1 + \hbar^n g$ , is equal to  $F + \hbar^n \chi$  mod  $\hbar^{n+1}$  and solves (5.3) mod  $\hbar^{n+2}$  since

$$\begin{aligned} (\Phi)_{\tilde{F}'} &= u^{\otimes 3} \cdot 1 \otimes \tilde{F} \cdot \text{id} \otimes \Delta(\tilde{F}) \cdot \text{id} \otimes \Delta(\Delta(u))^{-1} \cdot \Phi \\ &\quad \cdot \Delta \otimes \text{id}(\Delta(u)) \cdot \Delta \otimes \text{id}(\tilde{F}) \cdot \tilde{F} \otimes 1 \cdot (u^{\otimes 3})^{-1} \\ &= u^{\otimes 3} \cdot \Phi_D \cdot (u^{\otimes 3})^{-1} \\ &= \Phi_D \end{aligned}$$

where the first equality follows from the  $\mathfrak{g}$ -invariance of  $\Phi$  and the last from the  $\mathfrak{g}_D$ -invariance of  $u$  ■

We may therefore assume that  $\chi$  lies in  $(\bigwedge^2 \mathfrak{g})^{\mathfrak{l}_D}$ . We then note that, for  $n \geq 2$ ,

$$\begin{aligned} \xi(f; f_2, \dots, f_n + \chi) &= \xi(f; f_2, \dots, f_n) + f^{23}(\chi^{12} + \chi^{13}) + \chi^{23}(f^{12} + f^{13}) \\ &\quad - f^{12}(\chi^{13} + \chi^{23}) - \chi^{12}(f^{13} + f^{23}) \end{aligned}$$

so that  $F + \hbar^n \chi$  possesses an extension mod  $\hbar^{n+2}$  if, and only if,

$$\text{Alt}_3(\xi(f; f_2, \dots, f_n)) = \llbracket f, \chi \rrbracket$$

where we used lemma 5.7

**Proposition 5.10.** *The element  $\tilde{\xi} = \text{Alt}_3(\xi) \in (\bigwedge^3 \mathfrak{g})^{\mathfrak{l}_D}$  satisfies*

$$\llbracket f, \tilde{\xi} \rrbracket = 0 \quad \text{and} \quad \bar{\pi}_D^2(\tilde{\xi}) = 0$$

We defer the proof of proposition 5.10 to §5.6 in order to conclude the proof of theorem 5.2. By proposition 5.10,  $\tilde{\xi}$  is a 3-cocycle in  $((\bigwedge \mathfrak{g})^{\mathfrak{l}_D}, \llbracket f, \cdot \rrbracket)$  and we must show that it is a 3-coboundary. By theorem 4.5

$$\tilde{\xi} = \llbracket f, \chi \rrbracket + \eta \tag{5.14}$$

for some  $\chi \in (\bigwedge^2 \mathfrak{g})^{\mathfrak{l}_D}$  and

$$\eta \in (\bigwedge^3 \mathfrak{l}_D)^{\mathfrak{l}_D} = \bigwedge^3 \mathfrak{c}_D \oplus (\bigwedge^3 \mathfrak{g}_D)^{\mathfrak{g}_D} = (\bigwedge^3 \mathfrak{g}_D)^{\mathfrak{g}_D}$$

where the first equality follows from the fact that  $(\bigwedge^i \mathfrak{g}_D)^{\mathfrak{g}_D} = 0$  for  $i = 1, 2$  and the second from the assumption that  $|D_{\mathfrak{g}} \setminus D| \leq 2$  so that  $\mathfrak{c}_D$  is at most two-dimensional. Applying  $\bar{\pi}_D^3$  to both sides of (5.14), we find, since  $\bar{\pi}_D^2(f) = 0$  that

$$0 = \bar{\pi}_D^3(\llbracket f, \chi \rrbracket + \eta) = \eta$$

Note that if  $\Phi, F$  satisfy (5.4) and (5.5) respectively, then by lemma 5.5

$$\tilde{\xi}^{\ominus} = \text{Alt}_3(\xi^{\ominus}) = \text{Alt}_3(-\xi^{321}) = \tilde{\xi} \tag{5.15}$$

Since  $f^{\ominus} = -f$ , this implies

$$\llbracket f, \chi \rrbracket = \llbracket f, \chi \rrbracket^{\ominus} = -\llbracket f, \chi^{\ominus} \rrbracket$$

so that  $\chi' = 1/2(\chi - \chi^{\ominus})$  satisfies  $\chi'^{\ominus} = -\chi' = \chi'^{21}$  and  $\llbracket f, \chi' \rrbracket = \tilde{\xi}$  and  $F + \hbar^n \chi'$  is a solution of (5.3) mod  $\hbar^{n+1}$  possessing an extension to a solution mod  $\hbar^{n+2}$  satisfying (5.5). This completes the proof of theorem 5.2 ■

**Remark 5.11.** If  $\Phi$  satisfies (5.4), it is not necessary to ladder down, *i.e.*, assume that  $|D_{\mathfrak{g}} \setminus D| \leq 2$ . Indeed, by theorem 4.5, there exist

unique elements  $u \in \Lambda^3 \mathfrak{c}_D$ ,  $v \in (\Lambda^3 \mathfrak{g}_D)^{\mathfrak{g}^D}$  and a  $\chi \in (\Lambda^2 \mathfrak{g})^{l_D}$  such that

$$\tilde{\xi} = u + v + \llbracket f, \chi \rrbracket$$

Since  $\tilde{\xi}$  and  $u$  are killed by  $\bar{\pi}_D^3$ ,  $v = 0$ . Applying  $\Theta$  to both sides and using (5.15), we find that  $u^\Theta = u$ . This however implies that  $u = 0$  since  $\Theta$  acts by  $-1$  on  $\Lambda^3 \mathfrak{h} \supseteq \Lambda^3 \mathfrak{c}_D$ .

**5.6. Proof of proposition 5.10.** We begin with some preliminary lemmas. Let  $\tilde{\Delta} : U\mathfrak{g} \rightarrow U\mathfrak{g}^{\otimes 2}$  be a linear map. Let

$$\xi = \xi_1 \otimes \cdots \otimes \xi_k \in U\mathfrak{g}^{\otimes k}$$

$i \leq k$ , and write

$$\tilde{\Delta}(\xi_i) = \sum_a \xi'_{i,a} \otimes \xi''_{i,a}$$

For any enumeration  $j_1, \dots, j_{k+1}$  of  $[1, k+1]$ , we set

$$\zeta^{j_1, \dots, j_{i-1}, j_i, j_{i+1}, j_{i+2}, \dots, j_{k+1}} = \sum_a \eta_a$$

where  $\eta_a \in U\mathfrak{g}^{\otimes(k+1)}$  is the decomposable tensor with component  $\xi_\ell$  in position  $j_\ell$  if  $\ell \leq i-1$  and in position  $j_{\ell+1}$  if  $\ell \geq i+1$  and components  $\xi'_{i,a}, \xi''_{i,a}$  in positions  $j_i$  and  $j_{i+1}$  respectively. In other words,

$$\begin{aligned} \zeta^{j_1, \dots, j_{i-1}, j_i, j_{i+1}, j_{i+2}, \dots, j_{k+1}} &= \sigma \xi^{1, \dots, i-1, i, i+1, i+2, \dots, k+1} \\ &= \sigma \circ \text{id}^{\otimes(i-1)} \otimes \tilde{\Delta} \otimes \text{id}^{\otimes(k-i)} \xi \end{aligned}$$

where  $\sigma \in \mathfrak{S}_{k+1}$  is the permutation mapping  $\ell$  to  $i_\ell$ .

**Lemma 5.12.** *For any  $\xi \in \Lambda^k \mathfrak{g}$ , one has*

$$\begin{aligned} &(k+1) \text{Alt}_{k+1} \left( \sum_{i=1}^k (-1)^i \text{id}^{\otimes(i-1)} \otimes \tilde{\Delta} \otimes \text{id}^{\otimes(k-i)} \xi \right) \\ &= \sum_{1 \leq a < b \leq k+1} (-1)^{a+b} \left( (\text{Alt}_k \xi)^{ab, 1, \dots, \hat{a}, \dots, \hat{b}, \dots, k+1} - (\text{Alt}_k \xi)^{ba, 1, \dots, \hat{a}, \dots, \hat{b}, \dots, k+1} \right) \end{aligned} \tag{5.16}$$

**PROOF.** For any  $i \in [1, k]$  and  $a \neq b \in [1, k+1]$ , set

$$\mathfrak{S}_{k+1}^{i:a,b} = \{ \sigma \in \mathfrak{S}_{k+1} \mid \sigma(i) = a, \sigma(i+1) = b \}$$

Then, the left-hand side of (5.16) is equal to

$$\begin{aligned} & \frac{1}{k!} \sum_{\substack{1 \leq a < b \leq k+1 \\ 1 \leq i \leq k \\ \sigma \in \mathfrak{S}_{k+1}^{i:a,b}}} (-1)^i (-1)^\sigma \left( \xi^{\sigma(1), \dots, \sigma(i-1), ab, \sigma(i+2), \dots, \sigma(k+1)} \right. \\ & \quad \left. - \xi^{\sigma(1), \dots, \sigma(i-1), ba, \sigma(i+2), \dots, \sigma(k+1)} \right) \\ &= \frac{1}{k!} \sum_{\substack{1 \leq a < b \leq k+1 \\ 1 \leq i \leq k \\ \sigma \in \mathfrak{S}_{k+1}^{i:a,b}}} (-1)^i (-1)^\sigma \left( (\bar{\sigma}\xi)^{ab, 1, \dots, \hat{a}, \dots, \hat{b}, \dots, k+1} - (\bar{\sigma}\xi)^{ba, 1, \dots, \hat{a}, \dots, \hat{b}, \dots, k+1} \right) \end{aligned}$$

where, for any  $\sigma \in \mathfrak{S}_{k+1}^{i:a,b}$ ,

$$\bar{\sigma} \in \mathfrak{S}_k^i = \{\tau \in \mathfrak{S}_k \mid \tau(i) = 1\}$$

is the permutation determined by the commutativity of the following diagram

$$\begin{array}{ccc} [1, k] \setminus \{i\} & \longrightarrow & [1, k+1] \setminus \{i, i+1\} \\ \bar{\sigma} \downarrow & & \downarrow \sigma \\ [1, k] \setminus \{1\} & \longrightarrow & [1, k+1] \setminus \{a, b\} \end{array}$$

where the horizontal arrows are the obvious monotone identifications. Noting that  $\sigma \rightarrow \bar{\sigma}$  is an isomorphism of  $\mathfrak{S}_{k+1}^{i:a,b}$  onto  $\mathfrak{S}_k^i$  and deferring for the time being the proof that

$$(-1)^i (-1)^\sigma = (-1)^{a+b} (-1)^{\bar{\sigma}} \quad (5.17)$$

we see that the above is equal to

$$\frac{1}{k!} \sum_{\substack{1 \leq a < b \leq k+1 \\ 1 \leq i \leq k \\ \bar{\sigma} \in \mathfrak{S}_k^i}} (-1)^{a+b} (-1)^{\bar{\sigma}} \left( (\bar{\sigma}\xi)^{ab, 1, \dots, \hat{a}, \dots, \hat{b}, \dots, k+1} - (\bar{\sigma}\xi)^{ba, 1, \dots, \hat{a}, \dots, \hat{b}, \dots, k+1} \right)$$

and therefore to the right-hand side of (5.16). We turn now to the proof of (5.17). Let  $\bar{\bar{\sigma}} \in \mathfrak{S}_{k-1}$  be the permutation determined by the commutativity of

$$\begin{array}{ccccc} [1, k] \setminus \{i\} & \longrightarrow & [1, k-1] & \longrightarrow & [1, k+1] \setminus \{i, i+1\} \\ \bar{\sigma} \downarrow & & \bar{\bar{\sigma}} \downarrow & & \downarrow \sigma \\ [1, k] \setminus \{1\} & \longrightarrow & [1, k-1] & \longrightarrow & [1, k+1] \setminus \{a, b\} \end{array}$$

where the horizontal arrows are the obvious monotone identifications. Since  $(-1)^\sigma = (-1)^{\bar{\sigma}} \cdot (-1)^{i-1}$ , it suffices to prove that  $(-1)^\sigma = (-1)^{\bar{\sigma}}(-1)^{a+b-1}$ . This clearly holds if  $a = 1$  and  $b = 2$ . In the general case, letting  $\tau \in \mathfrak{S}_{k+1}$  be the unique permutation such that  $\tau$  is increasing on  $[1, k+1] \setminus \{a, b\}$ ,  $\tau(a) = 1$  and  $\tau(b) = 2$ , so that  $(-1)^\tau = (-1)^{a+b-1}$ , and noting that  $\overline{\tau \circ \sigma} = \bar{\sigma}$ , we see that

$$(-1)^\sigma = (-1)^{a+b-1}(-1)^{\tau \circ \sigma} = (-1)^{a+b-1}(-1)^{\overline{\tau \circ \sigma}} = (-1)^{a+b-1}(-1)^{\bar{\sigma}}$$

■

**Lemma 5.13.** *For any  $Y, X_1, \dots, X_k \in \mathfrak{g}$ , one has*

$$\begin{aligned} & Y \wedge X_1 \wedge \dots \wedge X_k \\ &= \frac{1}{(k+1)!} \sum_{\substack{1 \leq i \leq k+1 \\ \tau \in \mathfrak{S}_k}} (-1)^{i-1} (-1)^\tau X_{\tau(1)} \otimes \dots \otimes X_{\tau(i-1)} \otimes Y \otimes X_{\tau(i)} \otimes \dots \otimes X_{\tau(k)} \end{aligned} \quad (5.18)$$

PROOF. Set  $Z_1 = Y$  and  $Z_j = X_{j-1}$  for  $j = 2 \dots k+1$ . By definition,  $(k+1)!$  times the left-hand side of (5.18) is equal to

$$\begin{aligned} & \sum_{\tau \in \mathfrak{S}_{k+1}} (-1)^\tau Z_{\tau(1)} \otimes \dots \otimes Z_{\tau(k+1)} \\ &= \sum_{\substack{1 \leq j \leq k+1 \\ \tau \in \mathfrak{S}_{k+1}: \tau(j)=1}} (-1)^\tau X_{\tau(1)-1} \otimes \dots \otimes X_{\tau(j-1)-1} \otimes Y \otimes X_{\tau(j+1)-1} \otimes \dots \\ & \quad \dots \otimes X_{\tau(k+1)-1} \end{aligned}$$

For any  $\tau \in \mathfrak{S}_{k+1}$  such that  $\tau(j) = 1$ , let  $\bar{\tau} \in \mathfrak{S}_k$  be the permutation determined by the commutativity of

$$\begin{array}{ccc} [1, k] & \longrightarrow & [1, k+1] \setminus \{j\} \\ \bar{\tau} \downarrow & & \downarrow \tau \\ [1, k] & \longrightarrow & [1, k+1] \setminus \{1\} \end{array}$$

Then,  $(-1)^\tau = (-1)^{\bar{\tau}}(-1)^{j-1}$  and the above is equal to

$$\sum_{\substack{1 \leq j \leq k+1 \\ \tau \in \mathfrak{S}_{k+1}: \tau(j)=1}} (-1)^{j-1} (-1)^{\bar{\tau}} X_{\bar{\tau}(1)} \otimes \dots \otimes X_{\bar{\tau}(j-1)} \otimes Y \otimes X_{\bar{\tau}(j)} \otimes \dots \otimes X_{\bar{\tau}(k)}$$

which proves (5.18) ■

**Lemma 5.14.** *For any  $f \in \wedge^2 \mathfrak{g}$  and  $\eta \in \wedge^k \mathfrak{g}$ , one has*

$$\sum_{1 \leq a < b \leq k+1} (-1)^{a+b} [f^{ab}, \eta^{a,1,\dots,\widehat{a},\dots,\widehat{b},\dots,k+1} + \eta^{b,1,\dots,\widehat{a},\dots,\widehat{b},\dots,k+1}] = -\frac{k+1}{2} \llbracket f, \eta \rrbracket \quad (5.19)$$

PROOF. We may assume that  $f, \eta$  are of the form

$$\begin{aligned} f &= f_1 \wedge f_2 = \frac{1}{2}(f_1 \otimes f_2 - f_2 \otimes f_1) \\ \eta &= \eta_1 \wedge \dots \wedge \eta_k = \frac{1}{k!} \sum_{\sigma \in \mathfrak{S}_k} (-1)^\sigma \eta_{\sigma(1)} \otimes \dots \otimes \eta_{\sigma(k)} \end{aligned}$$

The left-hand side of (5.19) is then equal to

$$\begin{aligned} & \frac{1}{2k!} \sum_{\substack{1 \leq a < b \leq k+1 \\ \sigma \in \mathfrak{S}_k}} (-1)^{a+b} (-1)^\sigma \\ & \quad [f_1^a f_2^b - f_2^a f_1^b, (\eta_{\sigma(1)} \otimes \dots \otimes \eta_{\sigma(k)})^{a,1,\dots,\widehat{a},\dots,\widehat{b},\dots,k+1} \\ & \quad \quad \quad + (\eta_{\sigma(1)} \otimes \dots \otimes \eta_{\sigma(k)})^{b,1,\dots,\widehat{a},\dots,\widehat{b},\dots,k+1}] \\ &= \frac{1}{2k!} \sum_{\substack{1 \leq a < b \leq k+1 \\ \sigma \in \mathfrak{S}_k}} (-1)^{a+b} (-1)^\sigma \\ & \quad \eta_{\sigma(2)} \otimes \dots \otimes \eta_{\sigma(a)} \otimes [f_1, \eta_{\sigma(1)}] \otimes \eta_{\sigma(a+1)} \otimes \dots \\ & \quad \quad \quad \dots \otimes \eta_{\sigma(b-1)} \otimes f_2 \otimes \eta_{\sigma(b)} \otimes \dots \otimes \eta_{\sigma(k)} \\ & \quad - \eta_{\sigma(2)} \otimes \dots \otimes \eta_{\sigma(a)} \otimes f_2 \otimes \eta_{\sigma(a+1)} \otimes \dots \\ & \quad \quad \quad \dots \otimes \eta_{\sigma(b-1)} \otimes [f_1, \eta_{\sigma(1)}] \otimes \eta_{\sigma(b)} \otimes \dots \otimes \eta_{\sigma(k)} \\ & \quad - \eta_{\sigma(2)} \otimes \dots \otimes \eta_{\sigma(a)} \otimes [f_2, \eta_{\sigma(1)}] \otimes \eta_{\sigma(a+1)} \otimes \dots \\ & \quad \quad \quad \dots \otimes \eta_{\sigma(b-1)} \otimes f_1 \otimes \eta_{\sigma(b)} \otimes \dots \otimes \eta_{\sigma(k)} \\ & \quad + \eta_{\sigma(2)} \otimes \dots \otimes \eta_{\sigma(a)} \otimes f_1 \otimes \eta_{\sigma(a+1)} \otimes \dots \\ & \quad \quad \quad \dots \otimes \eta_{\sigma(b-1)} \otimes [f_2, \eta_{\sigma(1)}] \otimes \eta_{\sigma(b)} \otimes \dots \otimes \eta_{\sigma(k)} \end{aligned} \quad (5.20)$$

Setting  $\sigma' = \sigma \circ (1 \cdots a)$  in the first summand and  $\sigma' = \sigma \circ (1 \cdots b - 1)$  in the second, we see that their sum is equal to

$$\begin{aligned} & \frac{1}{2k!} \sum_{\substack{1 \leq a < b \leq k+1 \\ \sigma \in \mathfrak{S}_k}} (-1)^{b-1} (-1)^\sigma \\ & \quad \eta_{\sigma(1)} \otimes \cdots \otimes \eta_{\sigma(a-1)} \otimes [f_1, \eta_{\sigma(a)}] \otimes \eta_{\sigma(a+1)} \otimes \cdots \\ & \quad \quad \quad \cdots \otimes \eta_{\sigma(b-1)} \otimes f_2 \otimes \eta_{\sigma(b)} \otimes \cdots \otimes \eta_{\sigma(k)} \\ & - \frac{1}{2k!} \sum_{\substack{1 \leq a < b \leq k+1 \\ \sigma \in \mathfrak{S}_k}} (-1)^a (-1)^\sigma \\ & \quad \eta_{\sigma(1)} \otimes \cdots \otimes \eta_{\sigma(a-1)} \otimes f_2 \otimes \eta_{\sigma(a)} \otimes \cdots \\ & \quad \quad \quad \cdots \otimes \eta_{\sigma(b-2)} \otimes [f_1, \eta_{\sigma(b-1)}] \otimes \eta_{\sigma(b)} \otimes \cdots \otimes \eta_{\sigma(k)} \end{aligned}$$

and therefore to

$$\begin{aligned} & \frac{1}{2k!} \sum_{\substack{1 \leq a \neq b \leq k \\ \sigma \in \mathfrak{S}_k}} (-1)^{b-1} (-1)^\sigma \\ & \quad \eta_{\sigma(1)} \otimes \cdots \otimes \eta_{\sigma(a-1)} \otimes [f_1, \eta_{\sigma(a)}] \otimes \eta_{\sigma(a+1)} \otimes \cdots \\ & \quad \quad \quad \cdots \otimes \eta_{\sigma(b-1)} \otimes f_2 \otimes \eta_{\sigma(b)} \otimes \cdots \otimes \eta_{\sigma(k)} \\ & = \frac{k+1}{2} f_2 \wedge (\text{ad}(f_1) \eta) \end{aligned}$$

where we used lemma 5.13. Similarly, the sum of the last two summands in (5.20) is equal to

$$-\frac{k+1}{2} f_1 \wedge (\text{ad}(f_2) \eta)$$

Thus, the left-hand side of (5.19) is equal to

$$\frac{k+1}{2} (f_2 \wedge \text{ad}(f_1) \eta - f_1 \wedge \text{ad}(f_2) \eta) = -\frac{k+1}{2} \llbracket f_1 \wedge f_2, \eta \rrbracket$$

as claimed  $\blacksquare$

PROOF OF PROPOSITION 5.10. Write

$$(\Phi)_F = \Phi_D + \hbar^{n+1} \xi + \hbar^{n+2} \psi \quad \text{mod } \hbar^{n+3}$$

for some  $\psi \in U\mathfrak{g}^{\otimes 3}$ . Since  $\Phi_D$  is equal to 1 mod  $\hbar^2$  and satisfies the pentagon equation with respect to  $\Delta_F(\cdot) = F\Delta(\cdot)F^{-1}$ , we have, mod



$\hbar^{n+3}$ ,

$$\begin{aligned} 0 &= \text{Pent}_{\Delta_F}((\Phi)_F) \\ &= \text{Pent}_{\Delta_F}(\Phi_D) + \hbar^{n+1} d_H^{\Delta_F}(\xi) + \hbar^{n+2} d_H^{\Delta_F}(\psi) \\ &= \hbar^{n+1} d_H^{\Delta_F}(\xi) + \hbar^{n+2} d_H \psi \end{aligned}$$

where, for any  $\eta \in U\mathfrak{g}^{\otimes 3}$ ,

$d_H^{\Delta_F} \eta = 1 \otimes \eta - \Delta_F \otimes \text{id}^{\otimes 2}(\eta) + \text{id} \otimes \Delta_F \otimes \text{id}(\eta) - \text{id}^{\otimes 2} \otimes \Delta_F(\eta) + \eta \otimes 1$  is equal to  $d_H \text{ mod } \hbar$ . Applying  $\text{Alt}_4$  to both sides, and using lemmas 5.12 and 5.14, we find, with  $\tilde{\xi} = \text{Alt}_3 \xi \in (\bigwedge^3 \mathfrak{g})^{\text{td}}$

$$\begin{aligned} 0 &= \text{Alt}_4(d_H^{\Delta_F} \xi) \\ &= \frac{\hbar}{2} \sum_{1 \leq a < b \leq 4} (-1)^{a+b} [f^{ab}, \tilde{\xi}^{ab,1,\dots,\hat{a},\dots,\hat{b},\dots,4}] \\ &= \frac{\hbar}{2} \sum_{1 \leq a < b \leq 4} (-1)^{a+b} [f^{ab}, \tilde{\xi}^{a,1,\dots,\hat{a},\dots,\hat{b},\dots,4} + \tilde{\xi}^{b,1,\dots,\hat{a},\dots,\hat{b},\dots,4}] \\ &= -\hbar \llbracket f, \tilde{\xi} \rrbracket \end{aligned}$$

where we used the fact that, for any  $x \in U\mathfrak{g}$

$$\Delta_F(x) = \Delta(x) + \hbar[f, \Delta(x)] \quad \text{mod } \hbar^2$$

so that

$$\Delta_F(x) - \Delta_F(x)^{21} = 2\hbar[f, \Delta(x)]$$

This proves our claim since

$$\bar{\pi}_D^3 \tilde{\xi} = \text{Alt}_3 \bar{\pi}_D^3 \xi = 0$$

■

## 6. UNIQUENESS OF TWISTS

Let

$$\begin{aligned} \Phi &= 1 + \hbar^2 \varphi + \dots \in 1 + \hbar^2 (U\mathfrak{g}^{\otimes 3} \llbracket \hbar \rrbracket)^{\mathfrak{g}} \\ \Phi_D &= 1 + \hbar^2 \varphi_D + \dots \in 1 + \hbar^2 (U\mathfrak{g}_D^{\otimes 3} \llbracket \hbar \rrbracket)^{\mathfrak{g}^D} \end{aligned}$$

be two solutions of the pentagon equation (1.1) which are non-degenerate in the sense of definition 5.1. Contrary to §5, we do not assume in this subsection that  $\Phi_D = \bar{\pi}_D^3(\Phi)$  but merely that

$$\tilde{\varphi}_D = \bar{\pi}_D^3(\tilde{\varphi}) \tag{6.1}$$

where

$$\tilde{\varphi} = \text{Alt}_3 \varphi \in (\bigwedge^3 \mathfrak{g})^{\mathfrak{g}} \quad \text{and} \quad \tilde{\varphi}_D = \text{Alt}_3 \varphi_D \in (\bigwedge^3 \mathfrak{g}_D)^{\mathfrak{g}^D}$$

This implies in particular that if  $(\cdot, \cdot)$ ,  $(\cdot, \cdot)_D$  are the bilinear forms on  $\mathfrak{g}, \mathfrak{g}_D$  corresponding to  $\Phi, \Phi_D$  via (5.2) respectively, then  $(\cdot, \cdot)_D$  is the restriction of  $(\cdot, \cdot)$  to  $\mathfrak{g}_D$ . We denote the corresponding standard solutions of the MCYBE for  $\mathfrak{g}$  and  $\mathfrak{g}_D$  by  $r_{\mathfrak{g}}, r_{\mathfrak{g}_D}$ . Let now

$$F_i = 1^{\otimes 2} + \hbar f_i + \cdots \in 1 + \hbar(U\mathfrak{g}^{\otimes 2}[[\hbar]])^{\text{LD}}, \quad i = 1, 2$$

be two twists such that  $(\Phi)_{F_i} = \Phi_D$ . Since  $d_H f_i = 0$ , we have

$$\tilde{f}_i = \text{Alt}_2 f_i \in \left(\bigwedge^2 \mathfrak{g}\right)^{\text{LD}}$$

**Theorem 6.1.** *Let  $F_1, F_2$  be as above and assume that*

$$\tilde{f}_i = r_{\mathfrak{g}} - r_{\mathfrak{g}_D} + \nu_i$$

where  $\lambda_i \in \bigwedge^2 \mathfrak{c}_D$ . Then,

(i) *there exist elements*

$$u \in 1 + \hbar U\mathfrak{g}[[\hbar]]^{\text{LD}} \quad \text{and} \quad \lambda \in \hbar \bigwedge^2 \mathfrak{c}_D[[\hbar]]$$

such that

$$F_2 = \exp(\lambda) \cdot u \otimes u \cdot F_1 \cdot \Delta(u)^{-1} \quad (6.2)$$

(ii) *If*

$$\bar{\pi}_D^2(F_i) = 1, \quad i = 1, 2 \quad (6.3)$$

*u may be chosen such that  $\bar{\pi}_D^1(u) = 1$ .*

(iii) *If*

$$F_i^\Theta = F_i^{21}, \quad i = 1, 2 \quad (6.4)$$

*then  $\lambda = 0$  and u may be chosen such that  $u^\Theta = u$ . u is then unique with this property.*

(iv) *If  $|D_{\mathfrak{g}} \setminus D| \leq 1$ , then  $\lambda = 0$  and u is unique up to multiplication by  $\exp(c)$  for some  $c \in \hbar \mathfrak{c}_D[[\hbar]]$ .*

**PROOF.** (i)–(ii) Set

$$f = r_{\mathfrak{g}} - r_{\mathfrak{g}_D}$$

and write

$$f_i = d_H g_i + f + \nu_i$$

for some  $g_i \in U\mathfrak{g}^{\text{LD}}$ . Then, replacing  $F_i$  by

$$\exp(-\hbar \nu_i) \cdot (1 - \hbar g_i) \otimes (1 - \hbar g_i) \cdot F_i \cdot \Delta(1 - \hbar g_i)^{-1}$$

we may assume that

$$F_i = 1 + \hbar f \quad \text{mod } \hbar^2 \quad (6.5)$$

Note that if (6.3) holds, then, by corollary 2.5

$$0 = \bar{\pi}_D^2 f_i = d_H \bar{\pi}_D^1 g_i$$

and, replacing  $g_i$  by  $g_i - \bar{\pi}_D^2 g_i$ , we may assume that  $\bar{\pi}_D^1(g_i) = 0$ . Similarly, if (6.4) holds, then

$$d_H g_i^\Theta + f^\Theta + \nu_i^\Theta = d_H g_i + f^{21} + \nu_i^{21}$$

Since  $f^\Theta = -f = f^{21}$ , this yields

$$\nu_i^\Theta = -\nu_i \quad \text{and} \quad d_H g_i^\Theta = d_H g_i$$

whence  $\nu_i = 0$  since  $\Theta$  acts as multiplication by  $+1$  on  $\bigwedge^2 \mathfrak{h} \supseteq \bigwedge^2 \mathfrak{c}_D$  and, replacing  $g_i$  by  $1/2(g_i + g_i^\Theta)$ , we may assume that  $g_i^\Theta = g_i$ .

We wish now to construct two sequences

$$v_n \in U\mathfrak{g}^{\text{ld}} \quad \text{and} \quad \mu_n \in \bigwedge^2 \mathfrak{c}_D$$

such that, setting

$$u_n = (1 + \hbar^n v_n) \cdots (1 + \hbar v_1) \quad \text{and} \quad \lambda_n = \hbar^n \mu_n + \cdots + \hbar \mu_1$$

one has

$$F_2 = \exp(\lambda_n) \cdot u_n \otimes u_n \cdot F_1 \cdot \Delta(u_n)^{-1} \quad (6.6)$$

mod  $\hbar^{n+1}$ . If (6.3) (resp. (6.4)) holds, we require in addition that  $\bar{\pi}_D^1(v_n) = 0$  (resp.  $v_n^\Theta = v_n$  and  $\mu_n = 0$ ) for all  $n$ .

By (6.5), we may set  $v_1 = 0 = \mu_1$ . Assume therefore  $v_k, \mu_k$  constructed for  $k = 1 \dots n$  and some  $n \geq 1$ . Let  $F'_1$  be defined by the right-hand side of (6.6) so that

$$F_2 = F'_1 + \hbar^{n+1} \eta \quad \text{mod } \hbar^{n+2} \quad (6.7)$$

for some  $\eta \in (U\mathfrak{g}^{\otimes 2})^{\text{ld}}$ . One readily checks that  $(\Phi)_{F'_1} = \Phi_D$ . Subtracting from this the equation  $(\Phi)_{F_2} = \Phi_D$  and computing mod  $\hbar^{n+2}$ , we find that

$$d_H \eta = \eta^{23} + \text{id} \otimes \Delta(\eta) - \Delta \otimes \text{id}(\eta) - \eta^{12} = 0$$

Moreover,  $\bar{\pi}_D^2 \eta = 0$  (resp.  $\eta^\Theta = \eta^{21}$ ) if (6.3) (resp. (6.4)) holds. Thus,  $\eta = d_H v + \mu$  for some  $v \in U\mathfrak{g}^{\text{ld}}$  and  $\mu \in (\bigwedge^2 \mathfrak{g})^{\text{ld}}$  such that

$$\begin{aligned} \bar{\pi}_D^1 v &= 0, \\ v^\Theta &= v \quad \text{and} \quad \mu^\Theta = -\mu \end{aligned}$$

if (6.3), (6.4) hold respectively. Set  $v_{n+1} = -v$  and

$$\begin{aligned} F''_1 &= (1 + \hbar^{n+1} v_{n+1})^{\otimes 2} \cdot F'_1 \cdot \Delta(1 + \hbar^{n+1} v_{n+1})^{-1} \\ &= (1 + \hbar^{n+1} v_{n+1})^{\otimes 2} \cdot \exp(\lambda_n) \cdot u_n^{\otimes 2} \cdot F_1 \cdot \Delta(u_n)^{-1} \cdot \Delta(1 + \hbar^{n+1} v_{n+1})^{-1} \\ &= \exp(\lambda_n) \cdot ((1 + \hbar^{n+1} v_{n+1}) u_n)^{\otimes 2} \cdot F_1 \cdot \Delta((1 + \hbar^{n+1} v_{n+1}) u_n)^{-1} \end{aligned}$$

where the last equality stems from the fact that  $v_{n+1}$  is invariant under  $\mathfrak{l}_D$ . We have

$$F_2 = \exp(-\hbar^{n+1}\mu)F_1'' \quad \text{mod } \hbar^{n+2}$$

so the inductive step may be completed by setting  $\mu_{n+1} = -\mu$  provided we can show that  $\mu$  lies in  $\bigwedge^2 \mathfrak{c}_D$ . To see this, let

$$\overline{F}_2 = 1 + \hbar f + \hbar^2 f_2 + \cdots + \hbar^{n+1} f_{n+1}$$

be the truncation of  $F_2 \text{ mod } \hbar^{n+2}$  and define

$$\xi = \xi(f; f_2, \dots, f_{n+1}) \in (U\mathfrak{g}^{\otimes 3})^{\mathfrak{l}_D}$$

by

$$1 \otimes \overline{F}_2 \cdot \text{id} \otimes \Delta(\overline{F}_2) \cdot \Phi - \Phi_D \cdot \overline{F}_2 \otimes 1 \cdot \Delta \otimes \text{id}(\overline{F}_2) = \hbar^{n+2}\xi \quad \text{mod } \hbar^{n+3}$$

By lemma 5.4,  $d_H \xi = 0$  and, by corollary 5.6,  $\text{Alt}_3 \xi = 0$  since  $\overline{F}_2$  extends to a solution mod  $\hbar^{n+3}$ . Similarly, if  $\overline{F}_1''$  is the truncation of  $F_1'' \text{ mod } \hbar^{n+2}$ , the corresponding error  $\xi''$  satisfies  $d_H \xi'' = 0$  and  $\text{Alt}_3 \xi'' = 0$ . Since  $\overline{F}_1'' = \overline{F}_2 + \hbar^{n+1}\mu \text{ mod } \hbar^{n+2}$  and, for  $n \geq 1$

$$\begin{aligned} \xi'' - \xi &= \xi(f; f_2, \dots, f_{n+1} + \mu) - \xi(f; f_2, \dots, f_{n+1}) \\ &= f^{23}(\mu^{12} + \mu^{13}) + \mu^{23}(f^{12} + f^{13}) - f^{12}(\mu^{13} + \mu^{23}) - \mu^{12}(f^{13} + f^{23}) \end{aligned}$$

we find, using lemma 5.7, that  $\llbracket f, \mu \rrbracket = 0$ . By theorem 4.5, this implies that

$$\mu = \llbracket f, x \rrbracket + y$$

where  $y \in \bigwedge^2 \mathfrak{c}_D$  and  $x \in \mathfrak{g}^{\mathfrak{l}_D} = \mathfrak{c}_D \subseteq \mathfrak{h}$ . Since  $f$  is of weight 0,  $\llbracket f, x \rrbracket = -\text{ad}(x)f = 0$  whence  $\mu = y \in \bigwedge^2 \mathfrak{c}_D$ .

(iii) Let  $u \in 1 + \hbar(U\mathfrak{g}[[\hbar]])^{\mathfrak{l}_D}$ , with  $u^\Theta = u$ , be such that

$$u \otimes u \cdot F_1 \cdot \Delta(u)^{-1} = F_1 \quad (6.8)$$

We claim that  $u = 1$ . Assume that  $u = 1 \text{ mod } \hbar^n$  for some  $n \geq 1$  and write  $u = 1 + \hbar^n u_n \text{ mod } \hbar^{n+1}$ , where  $u_n \in U\mathfrak{g}^{\mathfrak{l}_D}$  is fixed by  $\Theta$ . Taking the coefficient of  $\hbar^{n+1}$  in (6.8) we find that  $d_H u_n = 0$ . This implies that  $u_n$  lies in  $\mathfrak{g}$  and therefore in  $\mathfrak{h}$  since it is of weight zero. Since  $\Theta$  acts as  $-1$  on  $\mathfrak{h}$  however,  $u_n = 0$  as claimed.

(iv) If  $|D_{\mathfrak{g}} \setminus D| \leq 1$ , then  $\bigwedge^2 \mathfrak{c}_D = 0$  so that  $\lambda = 0$ . Let now  $u \in 1 + \hbar(U\mathfrak{g}[[\hbar]])^{\mathfrak{l}_D}$  be such that

$$u \otimes u \cdot F_1 \cdot \Delta(u)^{-1} = F_1 \quad (6.9)$$

and write  $u = 1 + \hbar u_1 \text{ mod } \hbar^2$ . Taking the coefficient of  $\hbar$  in (6.9), we find that  $d_H u_1 = 0$  so that  $u_1 \in \mathfrak{g}^{\mathfrak{l}_D} = \mathfrak{c}_D$ . Now let  $u^{(2)} = u \cdot \exp(-\hbar u_1) = 1 + \hbar^2 u_2 \text{ mod } \hbar^2$ . Repeating the above argument with

$u^{(2)}$ , we find that  $u_2 \in \mathfrak{c}_D$  and finally that there exists a sequence  $u_n \in \mathfrak{c}_D$ ,  $n \geq 1$  such that

$$u = \prod_{n \geq 1} \exp(\hbar^n u_n) = \exp\left(\sum_{n \geq 1} \hbar^n u_n\right)$$

■

## REFERENCES

- [Di] J. Dixmier, *Algèbres enveloppantes*. Cahiers Scientifiques, Fasc. XXXVII. Gauthier–Villars Editeur, 1974.
- [DS] J. Donin, S. Shnider, *Cohomological construction of quantized universal enveloping algebras*, Trans. Amer. Math. Soc. **349** (1997), 1611–1632.
- [Dr1] V. G. Drinfeld, *Quantum groups*, Proceedings of the International Congress of Mathematicians (Berkeley 1986), 798–820, AMS, 1987.
- [Dr2] V. G. Drinfeld, *Quasi-Hopf algebras*, Leningrad Math. J. **1** (1990), 1419–1457.
- [EK] P. Etingof, D. Kazhdan, *Quantization of Lie bialgebras. I*, Selecta Math. (N.S.) **2** (1996), 1–41.
- [Lu] G. Lusztig, *Introduction to quantum groups*. Progress in Mathematics, 110. Birkhäuser, Boston, 1993.
- [MTL] J. J. Millson, V. Toledano Laredo, *Casimir operators and monodromy representations of generalised braid groups*, to appear in Transform. Groups **10** (2005), math.QA/0305062.
- [TL1] V. Toledano Laredo, *A Kohno–Drinfeld theorem for quantum Weyl groups*, Duke Math. J. **112** (2002), 421–451.
- [TL2] V. Toledano Laredo, *Flat connections and quantum groups*, The 2000 Twente Conference on Lie Groups. Acta Appl. Math. **73** (2002), 155–173.
- [TL3] V. Toledano Laredo, *Quasi-Coxeter algebras, Dynkin diagram cohomology and quantum Weyl groups*, math.QA/0506529.
- [TL4] V. Toledano Laredo, *Quasi-Coxeter quasitriangular quasibialgebras and the Casimir connection*, in preparation.

INSTITUT DE MATHÉMATIQUES DE JUSSIEU, UNIVERSITÉ PIERRE ET MARIE CURIE, UMR 7586, CASE 191, 16 RUE CLISSON, F-75013 PARIS  
*E-mail address:* toledano@math.jussieu.fr